

# Inconsistent centralised versus non-centralised ischaemic stroke aetiology

Yue Suo <sup>1,2</sup>, Jing Jing,<sup>1,2</sup> Xia Meng,<sup>1,2</sup> Zixiao Li <sup>1,2</sup>, Yuesong Pan <sup>1,2</sup>, Yong Jiang,<sup>1,2</sup> Xiaomeng Yang,<sup>1,2</sup> Huan Liu,<sup>1,2</sup> Hongyi Yan,<sup>1,2</sup> Liping Liu <sup>1,2</sup>, Xingquan Zhao,<sup>1,2</sup> Yilong Wang,<sup>1,2</sup> Hao Li <sup>1,2</sup>, Yongjun Wang <sup>1,2</sup>

**To cite:** Suo Y, Jing J, Meng X, et al. Inconsistent centralised versus non-centralised ischaemic stroke aetiology. *Stroke & Vascular Neurology* 2020;5:e000576. doi:10.1136/svn-2020-000576

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/svn-2020-000576>).

Received 10 August 2020

Revised 3 October 2020

Accepted 7 October 2020

Published Online First

5 November 2020

## ABSTRACT

**Background and purpose** The Trial of Org 10 172 in Acute Stroke Treatment (TOAST) system is the most widely used aetiological categorisation system in clinical practice and research. Limited studies have validated the accuracy of routine aetiological diagnosis of patients with ischaemic stroke according to the TOAST criteria when the reported subtype is assumed to be correct. We investigated the agreement between centralised and non-centralised (site-reported, at discharge) stroke subtypes in the Third China National Stroke Registry (CNSR-III), and analysed the influence of classification consistency on evaluation during hospitalisation and for secondary prevention strategy.

**Methods** All patients with ischaemic stroke from the CNSR-III study with complete diffusion-weighted imaging data were included. We used multivariable Cox proportional-hazard regression models to evaluate the factors associated with consistency between centralised and non-centralised stroke subtypes. Sensitivity analyses were conducted of the subgroup of patients with complete information.

**Results** This study included 12 180 patients (mean age, 62.3 years; and women, 31.4%). Agreement between centralised and non-centralised subtype was the highest for the large-artery atherosclerosis subtype stroke (77.4% of centralised patients), followed by the small-vessel occlusion subtype (40.6% of centralised patients). Agreements for cardioembolism and stroke of other determined aetiology subtypes were 38.7% and 12.2%, respectively. Patient-level and hospital-level factors were associated with the inconsistency between centralised/non-centralised aetiological subtyping. This inconsistency was related to differences in secondary prevention strategies. Only 15.3% of the newly diagnosed patients with cardioembolism underwent centralised subtyping with indications to receive oral anticoagulants at discharge. In comparison, 51.3% of the consistent cardioembolism group and 42.0% of the centrally reassigned cardioembolism group with anticoagulation indications were prescribed oral anticoagulants.

**Conclusions** Substantial inconsistency exists between centralised and non-centralised subtyping in China. Inaccurate aetiological subtyping could lead to inadequate secondary prevention, especially in patients with cardioembolic stroke.

## INTRODUCTION

Stroke has become the second leading cause of death worldwide<sup>1</sup> and the leading cause of mortality in China.<sup>2</sup> Ischaemic stroke

accounts for 69.6% of all strokes and the incidence of stroke has continued to steeply rise over the last three decades.<sup>3 4</sup> Although gradual improvements in stroke care have been achieved in recent years, regional differences still exist in the quality of diagnostic measures and stroke recurrence.<sup>5–7</sup>

Accurate classification of ischaemic stroke aetiology is indispensable for patient management and research because stroke prognosis<sup>8</sup> and strategies for secondary prevention of stroke differ by stroke subtype.<sup>9–11</sup> There are several aetiological subtyping systems<sup>12–19</sup>; however, despite the limited discriminatory ability and prognostic value,<sup>8</sup> the Trial of Org 10 172 in Acute Stroke Treatment (TOAST) system<sup>12</sup> remains the most widely used aetiological categorising system in clinical research and practice.<sup>20 21</sup>

In clinical practice and most clinical trials, aetiological diagnosis is based on the judgement of a local neurologist or site investigator. Previous studies have reported only moderate inter-rater reliability ( $\kappa$ : 0.42–0.64) with the TOAST system.<sup>22 23</sup> Both standardised medical record review<sup>24</sup> and centralised adjudication<sup>25</sup> have been shown to improve this situation. However, the accuracy of the routine aetiological diagnosis of ischaemic stroke according to TOAST criteria when the reported subtype is assumed to be correct has not been validated yet.

To the best of our knowledge, no large-scale study has compared the centralised versus non-centralised aetiological diagnosis of ischaemic stroke. The potential for heterogeneity in patient treatment following a centralised versus non-centralised diagnosis also warrants discussion. We, therefore, investigated the consistency between site-reported stroke subtypes at discharge according to centralised and non-centralised TOAST criteria for stroke diagnosis in the Third China National Stroke Registry (CNSR-III), its influence on aetiological evaluation during hospitalisation and its impact on secondary prevention strategy.



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Neurology, Beijing Tiantan Hospital, Beijing, China

<sup>2</sup>China National Clinical Research Center for Neurological Diseases, Beijing, China

## Correspondence to

Dr Yongjun Wang; [yongjunwang@ncrcnd.org.cn](mailto:yongjunwang@ncrcnd.org.cn)

## METHODS

### Study design

This study is an analysis of patients within the CNSR-III, a prospective patient cohort with ischaemic stroke or transient ischaemic attack (TIA) in China. Patients in our study were recruited between August 2015 and March 2018 from 201 study sites covering 26 provinces and municipalities in China.<sup>26</sup> Detailed inclusion and exclusion criteria are described in the CNSR-III study protocol.

### Study population and standardised diagnoses

The present analysis included all patients with ischaemic stroke with complete diffusion-weighted imaging (DWI) data as well as a complete aetiological evaluation and a non-centralised TOAST subtype designated by a site investigator at discharge. Site investigators and evaluators made decisions based on the 1993 version of the TOAST standard,<sup>12</sup> according to a previously published study protocol.<sup>26</sup> The TOAST system stratifies patients with ischaemic stroke into five subtypes: large artery atherosclerosis (LAA), cardioembolism (CE), small-vessel occlusion (SVO), stroke of other aetiology (OE) and stroke of undetermined cause (UE). A detailed description of the training process is included in the online supplemental materials (see online supplemental methods).

### Centralised and non-centralised subtyping

To ensure that the diagnosis standard was consistent, we adopted the use of just one version of the TOAST subtype system<sup>12</sup> for the centralised decision algorithm. A detailed methodology of the centralised and non-centralised subtyping is described in the supplementary methods (see online supplemental methods, online supplemental materials-screening report form and online supplemental materials-decision algorithm).

### Statistical analysis

A  $\chi^2$  test or a Fisher's exact test was used to compare categorical variables. Continuous variables were compared using either one-way analysis of variance or the Kruskal-Wallis test, where appropriate. The heterogeneity of centralised versus non-centralised subtype consistency was assessed using the  $\chi^2$  test. To analyse the potential causes of differences between centralised and non-centralised subtypes, inpatient department, hospital level and geographic location were also compared. Patients were further stratified into three categories per subtype based on centralised/non-centralised subtype consistency: consistent, centrally reassigned and newly diagnosed. For each specific subtype, the centrally reassigned group was defined as the group of patients diagnosed with the subtype at discharge but reassigned to a different subtype after centralised diagnosis. The newly diagnosed group for a given subtype included patients who were initially diagnosed with a subtype at discharge and then newly diagnosed with the given subtype after centralised diagnosis. For example, a patient originally diagnosed as OE and then centrally reassigned as CE would be included in

the centrally reassigned group for OE and the newly diagnosed group for CE. We used multivariate Cox proportional hazards regression to evaluate the factors associated with the inconsistency between centralised and non-centralised subtypes. Variables showing statistical differences in univariate analysis at baseline were modelled as confounding factors. We imputed missing values of the baseline variables used to adjust the regression models with mean or mode, as appropriate. No outcomes were imputed. In-hospital and discharge medication prescriptions were displayed for each category in every subtype. Indications of secondary prevention strategies were determined in accordance with established guidelines.<sup>10 27</sup> We performed sensitivity analyses using only the patient subgroup with complete information, that is, having DWI imaging,  $\geq 1$  extracranial artery evaluation,  $\geq 1$  intracranial artery evaluation,  $\geq 1$  cardiac rhythm evaluation (ECG (EKG) or 24-hour Holter) and  $\geq 1$  cardiac structure evaluation (transthoracic echocardiography (TTE) or transoesophageal echocardiography).

A two-sided *p* value  $< 0.05$  was considered to be statistically significant. All statistical analyses were performed using SAS V.9.4 (SAS Institute, Cary, North Carolina, USA).

## RESULTS


### Participant distribution and baseline characteristics

This study included a total of 12 180 patients with acute ischaemic stroke after excluding 1020 patients with TIA and 1966 patients with incomplete MRI data (figure 1). The included patients were older and had a lower rate of TIA history, myocardial infarction history and atrial fibrillation history; however, a higher percentage of patients had dyslipidaemia, were current smokers and or were diagnosed with cerebral infarction (see online supplemental table I).

Baseline characteristics of the study population are listed in online supplemental table II). The average age of the study population was 62.3 years (SD: 11.2 years), and 31.4% of all the included patients were women. A total of 11 586 (95.1%) patients underwent intracranial artery evaluation and 11 454 (94.0%) patients underwent a complete extracranial artery test. TTE data were available for 94.7% of the patients, EKG data for 94.3% and Holter data for 83.6%.

### Consistency between centralised and non-centralised subtype distributions

Differences between the distributions of aetiological subtypes based on discharge and those based on the centralised subtyping system are presented in figure 2. Substantial differences were observed between the centralised and non-centralised subtypes. Only 4316 patients (35.4%) were assigned to the same aetiological category after the centralised subtyping procedure performed by the site investigator at discharge. Agreement between the centralised and non-centralised subtypes was highest for



**Figure 1** Flow chart of patient inclusion and exclusion criteria.

the LAA subtype (77.4% of patients centrally diagnosed as LAA), followed by the SVO type (40.6% of patients centrally diagnosed as SVO). The agreement of CE and OE subtypes was 38.7% and 12.2%, respectively. Similar results were observed in the subgroup of patients with complete information.

To identify the potential reasons for the inconsistency between discharge and centralised subtype diagnosis, we compared both patient-level (demographic characteristics, baseline National Institutes of Health Stroke Scale (NIHSS) and so on) and hospital level (inpatient department, hospital type and geographic location or area of the hospital) factors (see online supplemental table III).


In multivariate logistic regression analysis, a history of ischaemic stroke (adjusted OR 0.841; 95% CI 0.749 to 0.943), TIA (adjusted OR 0.606; 95% CI 0.451 to 0.815) and dyslipidaemia (adjusted OR, 0.822; 95% CI 0.689 to 0.980) were each associated with fewer patients who were centrally reassigned from the LAA subtype. Additionally, hospitals in the urban area (adjusted OR 0.792; 95% CI 0.687 to 0.913) and tertiary hospitals (adjusted OR 0.748; 95% CI 0.627 to 0.891) were each associated with fewer patients centrally reassigned from the LAA subtype. Furthermore, we observed a significant interaction ( $p=0.005$ ) of hospital area with hospital type in the consistency of LAA diagnosis. Non-stroke unit inpatient departments were associated with more patients being centrally

reassigned from the LAA subtype (adjusted OR 1.140; 95% CI 1.012 to 1.284). Hospitals in the middle (adjusted OR 0.445; 95% CI 0.330 to 0.600) and the eastern (adjusted OR 0.476; 95% CI 0.359 to 0.631) regions of China were both associated with less newly diagnosed patients, indicating a more consistent LAA diagnosis.

A history of atrial fibrillation or flutter was associated with a more consistent diagnosis of the CE subtype (centrally reassigned CE group: adjusted OR 0.122; 95% CI 0.070 to 0.213; newly diagnosed CE group: adjusted OR 0.086; 95% CI 0.053 to 0.140). Evaluation of cardiac rhythm via 24-hour Holter was related to a more consistent CE subtype diagnosis (newly diagnosed CE group: adjusted OR 0.567; 95% CI 0.350 to 0.919).

A higher baseline NIHSS score was associated with more centrally reassigned patients with SVO (adjusted OR 1.360; 95% CI 1.164 to 1.588) and more newly diagnosed patients with SVO (adjusted OR 1.452; 95% CI 1.236 to 1.707). Additionally, tertiary hospitals were associated with more newly diagnosed patients with SVO (adjusted OR 1.410; 95% CI 1.140 to 1.743), while hospitals in urban areas were associated with fewer newly diagnosed patients with SVO (adjusted OR 0.446; 95% CI 0.360 to 0.552). An interaction was observed between hospital type, location/area and consistency of SVO diagnosis ( $p=0.037$ ).

Older age was associated with a high proportion of being centrally reassigned from the OE subtype (adjusted



**Figure 2** Centralised aetiological subtype and individualised subtype distribution. (A) Total number of each subtype in the study population. (B) Distribution of non-centralised subtyping results according to the percentages of centralised subtypes in the study population. (C) Total number of each subtype in the subgroup with complete information. (D) Distribution of non-centralised subtyping results according to the percentages of centralised subtypes in the subgroup with complete information. C, centralised; CE, cardioembolism; NC, non-centralised; OE, other determined cause; SVO, small-vessel occlusion; UE-multiple causes, with concomitant causes; UE-incomplete evaluation; UE-undetermined cause; LAA, large-artery atherosclerosis.

OR 1.062; 95% CI 1.022 to 1.102) and being newly diagnosed with OE (adjusted OR 1.062; 95% CI 1.016 to 1.109). Several patient-level and hospital-level factors were associated with agreement between centralised and non-centralised aetiological subtyping (table 1).

### Aetiological diagnosis and patient management

Fewer patients with newly diagnosed CE underwent 24-hour Holter (82.1%) than the patients in both the consistent CE (87.8%) and centrally reassigned CE

groups (87.7%; see online supplemental table III). The prescription rate of oral anticoagulants for secondary prevention was lower in newly diagnosed patients with CE (10% vs 50.2% for the consistent CE group and 27.6% for the centrally reassigned CE group,  $p < 0.001$ ). The prescription rate of antihypertensive agents was the highest in the consistent SVO group (48.7% vs 42.7% for the centrally reassigned SVO group and 44.3% for the newly diagnosed SVO group). Prescription rates

**Table 1** Multivariate analysis of potential causes of inconsistency between individualised aetiological classification and centralised classification of the study population

Characteristics		Centrally reassigned LAA		Newly diagnosed LAA	
Variables	Consistent LAA as reference	Adjusted OR (95% CI)	P value	Adjusted OR (95% CI)	P value
Medical history					
Ischaemic stroke	Yes	0.841 (0.749 to 0.943)	0.003	0.843 (0.687 to 1.035)	0.103
TIA	Yes	0.606 (0.451 to 0.815)	<0.001	0.585 (0.333 to 1.028)	0.062
Dyslipidaemia	Yes	0.822 (0.689 to 0.980)	0.029	0.806 (0.587 to 1.106)	0.181
Baseline NIHSS score	0–3	1.000 (ref)	Ref	1.000 (ref)	Ref
	≥4	0.731 (0.663 to 0.806)	<0.0001	0.537 (0.453 to 0.636)	<0.0001
Inpatient department	Stroke unit	1.000 (ref)	Ref	1.000 (ref)	Ref
	Other	1.140 (1.012 to 1.284)	0.031	1.037 (0.843 to 1.274)	0.733
Geographical region	West	1.000 (ref)	Ref	1.000 (ref)	Ref
	Middle	0.941 (0.770 to 1.150)	0.551	0.445 (0.330 to 0.600)	<0.0001
	East	0.882 (0.726 to 1.072)	0.207	0.476 (0.359 to 0.631)	<0.0001
Area	Rural	1.000 (ref)	Ref	1.000 (ref)	Ref
	Urban	0.792 (0.687 to 0.913)	0.001	1.163 (0.911 to 1.484)	0.226
Hospital type	Secondary	1.000 (ref)	Ref	1.000 (ref)	Ref
	Tertiary	0.748 (0.627 to 0.891)	0.001	0.796 (0.591 to 1.072)	0.133
P value of interaction for area and hospital type			0.005		0.214
Variables	Consistent CE as reference	Centrally reassigned CE Adjusted OR (95% CI)	P value	Newly diagnosed CE Adjusted OR (95% CI)	P value
Medical history					
Known atrial fibrillation or flutter	Yes	0.122 (0.070 to 0.213)	<0.0001	0.086 (0.053 to 0.140)	<0.0001
Diabetes mellitus	Yes	1.015 (0.599 to 1.721)	0.955	1.667 (1.060 to 2.623)	0.027
Baseline NIHSS score	0–3	1.000 (ref)	Ref	1.000 (ref)	Ref
	≥4	1.658 (1.131 to 2.432)	0.010	0.759 (0.541 to 1.065)	0.110
Evaluation of cardiac rhythm					
Holter	Yes	1.077 (0.606 to 1.915)	0.800	0.567 (0.350 to 0.919)	0.021
Variables	Consistent SVO as reference	Centrally reassigned SVO Adjusted OR (95% CI)	P value	Newly diagnosed SVO Adjusted OR (95% CI)	P value
Age		1.010 (1.003 to 1.018)	0.004	1.011 (1.003 to 1.019)	0.005
Sex		0.970 (0.813 to 1.158)	0.738	0.741 (0.613 to 0.896)	0.002
mRS prior to current event		1.072 (0.966 to 1.191)	0.192	1.215 (1.091 to 1.354)	<0.001
Medical history					
Current or previous smoker	Yes	0.780 (0.655 to 0.929)	0.005	0.858 (0.716 to 1.028)	0.096
Baseline NIHSS score	0–3	1.000 (ref)	Ref	1.000 (ref)	Ref
	≥4	1.360 (1.164 to 1.588)	0.0001	1.452 (1.236 to 1.707)	<0.0001
Geographical region	West	1.000 (ref)	Ref	1.000 (ref)	Ref
	Middle	1.320 (0.982 to 1.775)	0.066	1.447 (1.079 to 1.940)	0.014
	East	1.383 (1.039 to 1.839)	0.026	1.125 (0.849 to 1.490)	0.412
Area	Rural	1.000 (ref)	Ref	1.000 (ref)	Ref
	Urban	0.875 (0.713 to 1.074)	0.201	0.446 (0.360 to 0.552)	<0.0001
Hospital type	Secondary	1.000 (ref)	Ref	1.000 (ref)	Ref
	Tertiary	0.839 (0.662 to 1.064)	0.148	1.410 (1.140 to 1.743)	0.002

Continued

**Table 1** Continued

Characteristics		Centrally reassigned LAA		Newly diagnosed LAA	
Variables	Consistent LAA as reference	Adjusted OR (95% CI)	P value	Adjusted OR (95% CI)	P value
P value of interaction for area and hospital type			0.003		0.037
Variables	Consistent OE as reference	Centrally reassigned OE		Newly diagnosed OE	
		Adjusted OR (95% CI)	P value	Adjusted OR (95% CI)	P value
Age		1.062 (1.022 to 1.102)	0.002	1.062 (1.016 to 1.109)	0.007
Sex		0.293 (0.112 to 0.765)	0.012	0.701 (0.245 to 2.005)	0.507
Evaluation of cardiac rhythm					
EKG	Yes	1.702 (0.487 to 5.947)	0.405	10.173 (2.019 to 51.258)	0.005

West includes Inner Mongolia, Guangxi, Chongqing, Sichuan, Guizhou, Yunnan, Tibet, Shaanxi, Gansu, Qinghai, Ningxia and Xinjiang.

Middle includes Shanxi, Anhui, Jiangxi, Henan, Hubei and Hunan. East includes Beijing, Tianjin, Hebei, Shanghai, Jiangsu, Zhejiang, Fujian, Shandong, Guangdong, Hainan, Liaoning, Jilin and Heilongjiang.

CE, cardioembolism; LAA, large artery atherosclerosis; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; OE, other determined cause; ref, reference; SVO, small vessel occlusion.

for guideline-recommended treatments other than antithrombotic agents (such as lipid-lowering, antidiabetic and antihypertensive treatments) were comparable among the different groups based on the consistency of centralised and non-centralised subtyping in patients with LAA (table 2). Similar results were observed in the patient subgroup with complete information (see online supplemental table IV).

Of the 449 newly diagnosed patients with CE stroke, 431 (95.9%) had high-risk sources of CE. Among the high-risk sources, the largest proportion was newly diagnosed atrial fibrillation (58.47%), followed by a history of atrial fibrillation (28.54%; figure 3).

Oral anticoagulant prescription rates were significantly different among the three groups of patients with CE categorised by agreement between centralised and non-centralised subtyping. Among patients with indications for anticoagulation therapy, the prescription rate of oral anticoagulants at discharge in the newly diagnosed CE group was surprisingly lower (15.3%) than that in the consistent CE (51.3%) and the centrally reassigned CE groups (42.0%; table 3). Similar results were also observed in the patient subgroup with complete information (see online supplemental table V).

## DISCUSSION

In the current study, we found substantial differences between centralised and non-centralised aetiological classifications of stroke subtypes in China. Factors related to the heterogeneity between centralised and non-centralised subtyping included the hospital level, geographical region/location and area of the admitting hospital; the admitting department and stroke severity. Incomplete aetiological investigation during hospitalisation and inadequate secondary prevention strategies were observed in patients with inconsistent classification by site


evaluators (non-centralised) using centralised adjudication. Furthermore, we found that both a high baseline NIHSS score and admission to a stroke unit were each associated with a more consistent aetiological diagnosis between centralised and non-centralised stroke subtyping. Thus, differences in evaluation and care quality among patients with different severities might lead to inaccurate aetiological diagnoses in patients with mild neurological deficits.

The proportion of distribution of each subtype after centralised and non-centralised TOAST classification was similar to those reported in a previous registry,<sup>28</sup> with LAA accounting for a large proportion of ischaemic stroke and CE accounting for a small proportion. The particularly high prevalence of intracranial artery stenosis in Chinese patients with stroke<sup>29</sup> might explain the high-ranking proportion of the LAA subtype relative to the other subtypes. However, the proportion of LAA subtype among all patients classified by the non-centralised or centralised designations was different (60.8% vs 26.7%). A strict application of the TOAST classification criteria can lead to the designation of a significant number of strokes as an undetermined cause category.<sup>14</sup> A neurologist's 'clinical opinion', based on experience, might help to assign a high degree of confidence to one specific stroke cause in a particular patient when facing competing evidence of different aetiologies. Recent subtyping systems, such as the Chinese Ischaemic Stroke Subclassification (CISS) criteria, emphasise underlying pathological mechanisms and contain revised standards for categorising the LAA subtype.<sup>19</sup> According to CISS criteria, patients with infarct in the territory of an isolated penetrating artery and evidence of atherosclerotic plaque or any degree of stenosis in the parent artery are categorised into the LAA subtype, regardless of the degree of luminal stenosis of the clinically relevant artery. With the continuously

**Table 2** Patient medication based on consistency of centralised and discharge subtype

Treatment	Consistent LAA (n=2517)	Centrally reassigned LAA (n=4886)	Newly diagnosed LAA (n=735)	P value
Discharge treatment				
Antiplatelet	2302 (91.6%)	4483 (91.9%)	690 (93.9%)	0.134
Oral anticoagulation	6 (0.2%)	64 (1.3%)	8 (1.1%)	<0.001
Oral anticoagulation with indications*	0/1 to 0%	54/350 to 15.4%	0/0 to 0%	
Lipid-lowering in patients with dyslipidaemia	205/225 to 91.1%	344/360 to 95.6%	51/53 to 96.2%	
Antidiabetic in patients with diabetes	513/661 to 77.6%	917/1178 to 77.8%	135/169 to 79.9%	
Antihypertensive in patients with hypertension	979/1661 to 58.9%	1981/3121 to 63.5%	315/470 to 67.0%	
<b>Treatment</b>	<b>Consistent CE (n=284)</b>	<b>Centrally reassigned CE (n=266)</b>	<b>Newly diagnosed CE (n=449)</b>	<b>P value</b>
Discharge treatment				
Antiplatelet	121 (42.8%)	156 (59.8%)	361 (80.6%)	<0.001
Antiplatelet with indication within cardioembolic stroke†	2/10 20.0%	2/4 50.0%	5/6 83.3%	
Oral anticoagulation	142 (50.2%)	72 (27.6%)	45 (10.0%)	<0.001
Oral anticoagulation with indications*	139/272 to 51.1%	64/161 to 39.8%	44/301 to 14.6%	
Lipid-lowering in patients with dyslipidaemia	9/9 to 100%	14/14 to 100%	26/30 to 86.7%	
Antidiabetic in patients with diabetes	30/40 to 75.0%	35/46 to 76.1%	78/103 to 75.7%	
Antihypertensive in patients with hypertension	98/149 to 65.8%	77/143 to 53.8%	187/286 to 65.4%	
<b>Treatment</b>	<b>Consistent SVO (n=1193)</b>	<b>Centrally reassigned SVO (n=2003)</b>	<b>Newly diagnosed SVO (n=1749)</b>	<b>P value</b>
Discharge treatment				
Antiplatelet	1168 (97.9%)	1914 (95.6%)	1641 (93.9%)	<0.001
Oral anticoagulation	0 (0%)	15 (0.8%)	4 (0.2%)	0.002
Oral anticoagulation with indications*	0/0 to 0%	10/87 to 11.5%	0/0 to 0%	
Lipid-lowering in patients with dyslipidaemia	100/103 to 97.1%	133/138 to 96.4%	114/118 to 96.6%	
Antidiabetic in patients with diabetes	207/249 to 83.1%	361/431 to 83.8%	365/460 to 79.3%	
Antihypertensive in patients with hypertension	581/762 to 76.2%	855/1244 to 68.7%	775/1156 to 67.0%	
<b>Treatment</b>	<b>Consistent OE (n=20)</b>	<b>Centrally reassigned OE (n=451)</b>	<b>Newly diagnosed OE (n=144)</b>	<b>P value</b>
Discharge treatment				
Antiplatelet	17 (85%)	406 (90.6%)	122 (84.7%)	0.120
Oral anticoagulation	1 (5%)	7 (1.6%)	2 (1.4%)	0.478
Oral anticoagulation with indications*	0/0 to 0%	3/33 to 9.1%	0/0 to 0%	
Lipid-lowering in patients with dyslipidaemia	1/1 to 100%	32/33 to 97.0%	7/7 to 100%	
Antidiabetic in patients with diabetes	3/3 to 100%	81/110 to 73.6%	21/29 to 72.4%	
Antihypertensive in patients with hypertension	4/12 to 33.3%	197/282 to 69.9%	41/85 to 48.2%	

\*Indications of oral anticoagulation treatment included mechanical prosthetic valve, mitral stenosis with atrial fibrillation, left atrial/atrial appendage thrombus, left ventricular thrombus and atrial flutter. CE, cardioembolism; LAA, large artery atherosclerosis; OE, other determined cause; SVO, small vessel occlusion.



**Figure 3** Proportions of high-risk and medium-risk embolic sources for newly diagnosed patients with cardioembolic stroke. Panel A: proportions of high-risk sources. Panel B: proportions of medium-risk sources.

increasing development of novel techniques for plaque detection and vessel wall imaging, mild intracranial artery stenosis (<50% luminal stenosis) or non-stenotic atherosclerotic plaques are being recognised in >50% of patients with ischaemic symptoms.<sup>30</sup> These new subtyping criteria have had a profound influence on Chinese neurologists and might bias their ‘clinical opinions’ towards the LAA subtype.

Unlike the considerably high incidence of the CE subtype in the Caucasian population,<sup>31</sup> CE stroke accounts for only 5.7% of ischaemic strokes among Chinese patients, even with a complete aetiological investigation. Disagreements between non-centralised and centralised aetiological diagnosis of the CE subtype were mainly caused by the detection of atrial fibrillation (AF) after ischaemic stroke. AF is a well-established predictor of stroke recurrence and has a validated association with cardiogenic embolism.<sup>32</sup> The AF-SCREEN International Collaboration recommends 72-hour electrocardiographic monitoring for AF detection.<sup>33</sup> Because the CNSR-III protocol was written before publication of the white paper,<sup>33</sup> a stepwise approach was used to evaluate AF using resting EKG followed by 24-hour Holter monitoring. The current diagnostic approach detected AF-related stroke in 252 patients and reassigned them to the CE category. Another reason for subtyping inconsistency might be the availability of complementary information from the centralised data collection. Newly observed high-risk or medium-risk cardiac embolism sources within centralised data contributed to 62.3% of cardioembolic stroke classifications.

Inconsistency between non-centralised and centralised classifications of the CE subtype was associated with

inadequate guideline-recommended secondary prevention strategy, especially with regard to indications for treatment with oral anticoagulants. Among patients with ischaemic stroke with non-valvular atrial fibrillation (NVAf) in the CNSR without contraindications to anticoagulation, only 16.2% received warfarin therapy as secondary prevention.<sup>34</sup> In the second CNSR (CNSR-II), the prescription rate of warfarin for patients with stroke with NVAf remained low (19.4%).<sup>35</sup> Although an overall increase was observed in the use of oral anticoagulants in the current analysis, we found that the prescription rate in patients with unrecognised CE with indications of anticoagulation treatment (15.3%) was even lower than that reported in previous studies.<sup>34,35</sup> Inaccurate diagnosis of CE aetiology could cause suboptimal anticoagulation treatment in patients with CE indicated for anticoagulant therapy. Enhancing the accuracy and reliability of aetiological diagnosis is an essential step to improve adherence to guideline-recommended secondary prevention treatment.

Our study has several strengths. We used an algorithm for centralised adjudication of aetiology categories, which integrated all necessary data elements collected and processed centrally by trained specialists. Furthermore, this study was based on a cohort in which >94% of all included patients underwent complete aetiological examinations. The centralised aetiology classification was reproducible and stable because it eliminated any disagreements between evaluators in the judgement step.

However, this study had certain limitations. First, the algorithm was designed based on previously published rules.<sup>12</sup> New perspectives to stratify the sources of cardiac embolism risk are continually being updated.<sup>36</sup>

**Table 3** Oral anticoagulant treatment of patients with cardioembolic stroke with unchanged or reassigned subtype

Treatment	Consistent CE* (n=282)				Centrally reassigned CE† (n=262)				Newly diagnosed CE‡ (n=447)			
	With indication§		Without indication		With indication§		Without indication		With indication§		Without indication	
	n=270	P value	n=12	P value	n=161	P value	n=101	P value	n=300	P value	n=147	P value
In-hospital anticoagulant	151 (55.9%)	0.146	4 (33.3%)	0.146	86 (53.4%)	0.146	18 (17.8%)	<0.0001	82 (27.3%)	<0.0001	12 (8.2%)	<0.0001
Warfarin	89/151 to 58.9%	0.647	3/4 to 75.0%	0.647	46/86 to 53.5%	0.647	5/18 to 27.8%	0.069	26/82 to 31.7%	0.069	1/12 to 8.3%	0.169
LWHP	80/151 to 53.0%	0.349	1/4 to 25.0%	0.349	49/86 to 57.0%	0.349	9/18 to 50.0%	0.611	43/82 to 52.4%	0.611	8/12 to 66.7%	0.537
Heparin	1/151 to 0.7%	1.000	0/4 to 0%	1.000	4/18 to 22.2%	1.000	0/86 to 0%	<0.001	1/82 to 1.2%	<0.001	0/12 to 0%	1.000
Rivaroxaban	9/151 to 6.0%	1.000	0/4 to 0%	1.000	2/86 to 2.3%	1.000	0/18 to 0%	1.000	2/82 to 2.4%	1.000	0/12 to 0%	1.000
Dabigatran	22/151 to 14.6%	0.478	1/4 to 25.0%	0.478	6/86 to 7.0%	1.000	1/18 to 5.6%	1.000	5/82 to 6.1%	1.000	0/12 to 0%	1.000
Apixaban	0/151 to 0%	NE	0/4 to 0%	NE	0/86 to 0%	NE	0/18 to 0%	NE	0/82 to 0%	NE	0/12 to 0%	NE
Others	4/151 to 2.7%	1.000	0/4 to 0%	1.000	2/86 to 2.3%	1.000	1/18 to 5.6%	0.438	11/82 to 13.4%	0.438	3/12 to 25.0%	0.379
<b>Newly diagnosed CE†† (n=448)</b>												
Discharge anticoagulant	139 (51.3%)	0.252	4 (33.3%)	0.252	66 (42.0%)	0.252	8 (7.7%)	<0.0001	46 (15.3%)	<0.0001	2 (1.4%)	<0.0001
Warfarin	103/139 to 74.1%	1.000	3/4 to 75.0%	1.000	53/66 to 80.3%	1.000	7/8 to 87.5%	1.000	33/46 to 71.7%	1.000	1/2 to 50.0%	0.503
LWHP	0/139 to 0%	NE	0/4 to 0%	NE	2/66 to 3.0%	NE	1/8 to 12.5%	0.294	2/46 to 4.4%	0.294	1/2 to 50.0%	0.122
Heparin	0/139 to 0%	NE	0/4 to 0%	NE	0/66 to 0%	NE	0/8 to 0%	NE	0/46 to 0%	NE	0/2 to 0%	NE
Rivaroxaban	8/139 to 5.8%	1.000	0/4 to 0%	1.000	4/66 to 6.1%	1.000	1/8 to 12.5%	0.445	4/46 to 8.7%	0.445	0/2 to 0%	1.000
Dabigatran	27/139 to 19.4%	1.000	1/4 to 25.0%	1.000	7/66 to 10.6%	1.000	0/8 to 0%	1.000	7/46 to 15.2%	1.000	0/2 to 0%	1.000
Apixaban	0/139 to 0%	NE	0/4 to 0%	NE	0/66 to 0%	NE	0/8 to 0%	NE	0/46 to 0%	NE	0/2 to 0%	NE
Others	1/139 to 0.7%	1.000	0/4 to 0%	1.000	0/66 to 0%	1.000	0/8 to 0%	NE	0/46 to 0%	NE	0/2 to 0%	NE

\*Missing data of two patients.

†Missing data of four patients.

‡Missing data of two patients.

§Indications of oral anticoagulation treatment included mechanical prosthetic valve, mitral stenosis with atrial fibrillation, atrial fibrillation, left atrial/atrial appendage thrombus, left ventricular thrombus and atrial flutter.

¶Missing data of one patient.

\*\*Missing data of five patients.

††Missing data of one patient.

CE, cardioembolism; LWHP, low molecular weight heparin; NE, not estimable.

Efficient treatments for different potential embolic sources continue to be developed, such as percutaneous patent foramen ovale (PFO) closure for PFO-related stroke.<sup>37–39</sup> For future use, the algorithm employed in the current study needs to be updated with the most recent knowledge. Second, many patients in the current study were assigned to the undetermined causes category with  $\geq 1$  competing mechanism after completing the diagnostic workup. It is thus essential to refine the hierarchical competing standard among subtype elements. Third, the standard evaluation process in the current study lacked cardiac MRI, 72-hour EKG monitoring and specific biomarkers for cardiac function. Fourth, the evaluators at each study site were trained in the use of videotape. The intra-rater and inter-rater reliabilities were not centrally assessed before aetiological diagnosis was performed. However, this reflects a real-world challenge of evaluating interobserver and intraobserver agreements of aetiological diagnoses owing to the large scale of neurological departments and constantly rotating physicians in the many hospitals in China. Efforts to improve the quality of centralised training processes and constant interactive feedback are needed to narrow the gap between non-centralised and centralised subtyping in future multi-centre studies.

## CONCLUSIONS

We report substantial differences between discharge and centralised diagnosis of stroke aetiology. The factor most associated with these differences was the inhomogeneity among admitting departments and hospitals in their ability to identify potential stroke mechanisms. An inadequate secondary prevention strategy was observed in patients with inconsistent classification by site evaluators (non-centralised) and centralised adjudicators.

**Twitter** Yilong Wang @yilong

**Contributors** Y-JW had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Y-JW, JJ and YS designed the protocol and framework of this study. YS, JJ, XM, YJ and YP collected the data. Huan Liu and HY conducted statistical analysis. YS drafted the manuscript. ZL, YJ, XY, LL, XZ, Y-LW and Hao Li reviewed, edited and approved the final version.

**Funding** This study was funded by National Science and Technology Major Project National Key R&D Program of China Beijing Municipal Science & Technology Commission (D171100003017002).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Yue Suo <http://orcid.org/0000-0002-0632-0262>

Zixiao Li <http://orcid.org/0000-0002-4713-5418>

Yuesong Pan <http://orcid.org/0000-0003-3082-6789>

Liping Liu <http://orcid.org/0000-0003-2943-055X>

Hao Li <http://orcid.org/0000-0002-8591-4105>

Yongjun Wang <http://orcid.org/0000-0002-9976-2341>

## REFERENCES

- GBD 2016 Stroke Collaborators. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the global burden of disease study 2016. *Lancet Neurol* 2019;18:439–58.
- Zhou M, Wang H, Zeng X, et al. Mortality, morbidity, and risk factors in China and its provinces, 1990–2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2019;394:1145–58.
- Wang W, Jiang B, Sun H, et al. Prevalence, Incidence, and Mortality of Stroke in China: Results from a Nationwide Population-Based Survey of 480 687 Adults. *Circulation* 2017;135:759–71.
- Global Burden of Disease Collaborative Network. Global burden of disease study 2017 (GBD 2017) results, 2018. Available: <http://ghdx.healthdata.org/gbd-results-tool> [Accessed Aug 2020].
- Gu H-Q, Yang X, Rao Z-Z, et al. Disparities in outcomes associated with rural-urban insurance status in China among inpatient women with stroke: a registry-based cohort study. *Ann Transl Med* 2019;7:426.
- Li Z, Jiang Y, Li H, et al. China's response to the rising stroke burden. *BMJ* 2019;364:l879.
- Wang Y, Li Z, Wang Y, et al. Chinese stroke center alliance: a national effort to improve healthcare quality for acute stroke and transient ischaemic attack: rationale, design and preliminary findings. *Stroke Vasc Neurol* 2018;3:256–62.
- Arsava EM, Helenius J, Avery R, et al. Assessment of the predictive validity of etiologic stroke classification. *JAMA Neurol* 2017;74:419–26.
- Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American heart Association/American stroke association. *Stroke* 2018;2018:e46–110.
- Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American heart Association/American stroke association. *Stroke* 2014;45:2160–236.
- National Collaborating Centre for Chronic Conditions. *National clinical guideline for diagnosis and initial management of acute stroke and transient ischaemic attack (TIA)*. London, United Kingdom: Royal College of Physicians of London, 2008.
- Adams HP, Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute ischemic stroke. definitions for use in a multicenter clinical trial. TOAST. trial of ORG 10172 in acute stroke treatment. *Stroke* 1993;24:35–41.
- Ay H, Benner T, Arsava EM, et al. A computerized algorithm for etiologic classification of ischemic stroke: the causative classification of stroke system. *Stroke* 2007;38:2979–84.
- Ay H, Furie KL, Singhal A, et al. An evidence-based causative classification system for acute ischemic stroke. *Ann Neurol* 2005;58:688–97.
- Arsava EM, Ballabio E, Benner T, et al. The causative classification of stroke system: an international reliability and optimization study. *Neurology* 2010;75:1277–84.
- Amarenco P, Bogousslavsky J, Caplan LR, et al. New approach to stroke subtyping: the A-S-C-O (phenotypic) classification of stroke. *Cerebrovasc Dis* 2009;27:502–8.
- Amarenco P, Bogousslavsky J, Caplan LR, et al. The ASCOD phenotyping of ischemic stroke (updated ASCO phenotyping). *Cerebrovasc Dis* 2013;36:1–5.
- Chen P-H, Gao S, Wang Y-J, et al. Classifying ischemic stroke, from TOAST to CISS. *CNS Neurosci Ther* 2012;18:452–6.
- Gao S, Wang YJ, Xu AD, et al. Chinese ischemic stroke subclassification. *Front Neurol* 2011;2:6.
- Perera KS, KKH N, Nayar S, et al. Association between low-dose rivaroxaban with or without aspirin and ischemic stroke subtypes: a secondary analysis of the COMPASS trial. *JAMA neurol* 2019;77:43–8.
- Mazzucco S, Li L, Binney L, et al. Prevalence of patent foramen ovale in cryptogenic transient ischaemic attack and non-disabling stroke at older ages: a population-based study, systematic review, and meta-analysis. *Lancet Neurol* 2018;17:609–17.
- Atiya M, Kurth T, Berger K, et al. Interobserver agreement in the classification of stroke in the women's health study. *Stroke* 2003;34:565–7.
- Gordon DL, Bendixen BH, Adams HP, et al. Interphysician agreement in the diagnosis of subtypes of acute ischemic stroke:

- implications for clinical trials. The TOAST Investigators. *Neurology* 1993;43:1021–7.
- 24 Goldstein LB, Jones MR, Matchar DB, *et al.* Improving the reliability of stroke subgroup classification using the trial of ORG 10172 in acute stroke treatment (TOAST) criteria. *Stroke* 2001;32:1091–7.
  - 25 Majersik JJ, Cole JW, Golledge J, *et al.* Recommendations from the International stroke genetics Consortium, part 1: standardized phenotypic data collection. *Stroke* 2015;46:279–84.
  - 26 Wang Y, Jing J, Meng X, *et al.* The third China national stroke registry (CNSR-III) for patients with acute ischaemic stroke or transient ischaemic attack: design, rationale and baseline patient characteristics. *Stroke Vasc Neurol* 2019;4:158–64.
  - 27 Liu L, Chen W, Zhou H, *et al.* Chinese stroke association guidelines for clinical management of cerebrovascular disorders: Executive summary and 2019 update of clinical management of ischaemic cerebrovascular diseases. *Stroke Vasc Neurol* 2020;5:159–76.
  - 28 Wang Y, Xu J, Zhao X, *et al.* Association of hypertension with stroke recurrence depends on ischemic stroke subtype. *Stroke* 2013;44:1232–7.
  - 29 Wang Y, Zhao X, Liu L, *et al.* Prevalence and outcomes of symptomatic intracranial large artery stenoses and occlusions in China: the Chinese intracranial atherosclerosis (CICAS) study. *Stroke* 2014;45:663–9.
  - 30 Wang Y, Liu X, Wu X, *et al.* Culprit intracranial plaque without substantial stenosis in acute ischemic stroke on vessel wall MRI: a systematic review. *Atherosclerosis* 2019;287:112–21.
  - 31 Ornello R, Degan D, Tiseo C, *et al.* Distribution and temporal trends from 1993 to 2015 of ischemic stroke subtypes: a systematic review and meta-analysis. *Stroke* 2018;49:814–9.
  - 32 Lip GYH, Hunter TD, Quiroz ME, *et al.* Atrial fibrillation diagnosis timing, ambulatory ECG monitoring utilization, and risk of recurrent stroke. *Circ Cardiovasc Qual Outcomes* 2017;10:e002864.
  - 33 Schnabel RB, Haeusler KG, Healey JS, *et al.* Searching for atrial fibrillation poststroke: a white paper of the AF-SCREEN international collaboration. *Circulation* 2019;140:1834–50.
  - 34 Wang C, Yang Z, Wang C, *et al.* Significant underuse of warfarin in patients with nonvalvular atrial fibrillation: results from the China national stroke Registry. *J Stroke Cerebrovasc Dis* 2014;23:1157–63.
  - 35 Yang X, Li Z, Zhao X, *et al.* Use of warfarin at discharge among acute ischemic stroke patients with nonvalvular atrial fibrillation in China. *Stroke* 2016;47:464–70.
  - 36 Pepi M, Evangelista A, Nihoyannopoulos P, *et al.* Recommendations for echocardiography use in the diagnosis and management of cardiac sources of embolism: European association of echocardiography (EAE) (a registered branch of the ESC). *Eur J Echocardiogr* 2010;11:461–76.
  - 37 Saver JL, Carroll JD, Thaler DE, *et al.* Long-Term outcomes of patent foramen ovale closure or medical therapy after stroke. *N Engl J Med* 2017;377:1022–32.
  - 38 Mas J-L, Derumeaux G, Guillon B, *et al.* Patent foramen ovale closure or anticoagulation vs. antiplatelets after stroke. *N Engl J Med* 2017;377:1011–21.
  - 39 Søndergaard L, Kasner SE, Rhodes JF, *et al.* Patent foramen ovale closure or antiplatelet therapy for cryptogenic stroke. *N Engl J Med* 2017;377:1033–42.