

H₂S promotes developmental brain angiogenesis via the NOS/NO pathway in zebrafish

Weiying Jiang¹,^{ORCID} Chen Liu,² Mingzhu Deng,² Fei Wang,¹ Xiao Ren,¹ Yilin Fan,¹ Jiulin Du,³ Yonggang Wang¹

To cite: Jiang W, Liu C, Deng M, et al. H₂S promotes developmental brain angiogenesis via the NOS/NO pathway in zebrafish. *Stroke & Vascular Neurology* 2021;6:e000584. doi:10.1136/svn-2020-000584

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/svn-2020-000584>).

Received 15 August 2020
Revised 19 October 2020
Accepted 28 October 2020
Published Online First
27 November 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Neurology, Shanghai Jiao Tong University School of Medicine Affiliated Renji Hospital, Shanghai, China
²Department of Neurology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China
³Institute of Neuroscience, State Key Laboratory of Neuroscience, Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai, China

Correspondence to

Dr Yonggang Wang;
w100yg@163.com

ABSTRACT

Background Hydrogen sulphide (H₂S) is considered as the third member of the gasotransmitter family, along with nitric oxide (NO) and carbon monoxide. H₂S has been reported to induce angiogenesis by promoting the growth, migration and tube-like structure formation of endothelial cells. Those studies were conducted in conditions of cell culture, mouse Matrigel plug assay model, rat wound healing model or rat hindlimb ischaemia model. Recent in vivo studies showed the physiological importance of H₂S in muscle angiogenesis. However, the importance of endogenous H₂S for brain angiogenesis during development remains unknown. We therefore aimed at determining the role of H₂S in brain vascular development.

Methods and results Both knockdown and knockout of H₂S-producing enzymes, cystathionine β-synthase (*cbs*) and cystathionine γ-lyase (*cth*), using morpholino oligonucleotides and clustered regularly interspaced short palindromic repeats/Cas9-mediated mutation, impaired brain vascular development of larval zebrafish. Incubation with the slow-releasing H₂S donor GYY4137 alleviated the defects of brain vascular development in *cbs* and *cth* morphants. Quantitative analysis of the midbrain vascular network showed that H₂S enhances angiogenesis without affecting the topological structure of the brain vasculature. Mechanically, nitric oxide synthase 2a (*nos2a*) expression and NO production were decreased in both *cbs* and *cth* morphants. Overexpression of *nos2a* by coinjection of *cbs* or *cth* MO with full-length zebrafish *nos2a* mRNA alleviated the brain vascular developmental defects in *cbs* and *cth* morphants.

Conclusion We conclude that H₂S promotes brain developmental angiogenesis via the NOS/NO pathway in zebrafish.

INTRODUCTION

Since found in the mammalian brain in 1989,¹ hydrogen sulphide (H₂S) has been attracting continuous attention on its physiological and pathophysiological roles in the central nervous system (CNS). Along with nitric oxide (NO) and carbon monoxide, H₂S has been considered to be the third member of the gasotransmitter family.² Cystathionine β-synthase (CBS) and cystathionine γ-lyase (CTH) are two main H₂S-producing enzymes from cysteine and homocysteine in mammalian tissues.³ CBS is abundantly expressed in

the brain,⁴ especially in the hippocampus and the cerebellum.⁵ In developing mouse brains, H₂S is preferentially produced in radial glia/astrocyte lineages.^{6,7} The expression of CTH is rather widely distributed among peripheral tissues including the liver, pancreas, uterus and intestine.⁸ Western blotting experiments detected CTH rather than CBS in cerebral microvessels of newborn pigs⁹ and CTH is the main H₂S-producing enzyme in blood vessels.¹⁰

In the CNS, the well-studied functions of H₂S include the modulation of neurotransmission⁴ and neuroprotection.¹¹ In the peripheral vascular system, H₂S is reported to induce vasorelaxation¹² and angiogenesis.^{13–15} In vitro studies have demonstrated that exogenous H₂S at physiologically relevant doses induced angiogenesis by promoting the growth, migration and tube-like structure formation of vascular endothelial cells (ECs),^{13,14} while in situ studies showed that neovascularisation is promoted by H₂S in the mouse Matrigel plug assay model,¹³ rat wound healing model¹⁴ and rat hindlimb ischaemia model.¹⁵ Recently, two in vivo studies have shown the importance of H₂S in muscle angiogenesis.^{16,17} However, the importance of endogenous H₂S for brain angiogenesis during development remains unknown.

The zebrafish is a well-established vertebrate model for in vivo study of vascular development.¹⁸ Its accessible, small and transparent embryo facilitates high resolution in vivo imaging of the brain vasculature.¹⁹ In the present study, we used larval zebrafish as a model and found that endogenous H₂S plays an important role in the brain vascular development by promoting angiogenesis, and this effect is mainly mediated through the nitric oxide synthase (NOS)/NO pathway.

METHODS

Detailed methods are available in the online supplemental material.

In vivo confocal imaging

In vivo long-term serial confocal imaging of the midbrain vasculature was conducted in the same larva during 3–5 days post fertilisation (dpf). Larvae were embedded in 1.2% low-melting agarose (Invitrogen, 16520050) without anaesthetic at room temperature. After being imaged, the 3 or 4-dpf larvae were dug out for husbandry till subsequent imaging of the same larva was performed. Imaging was carried out with an Olympus Fluoview 1000 confocal microscope (Olympus, Japan). XLumplfl 20× (W/IR; NA, 0.95) objective lenses were used. The Z-step of imaging was 3 µm. Raw images were processed with ImageJ. Quantitative analysis of the midbrain vasculature morphology was conducted following the processes previously described.¹⁹

Whole-mount in situ hybridization

Zebrafish whole-mount in situ hybridization was performed as previously described.^{20, 21} Probes for the *cbs* and *cth* mRNAs were synthesised and labelled with digoxigenin (Roche, Mannheim, Germany). Embryos were incubated with corresponding probes (1 ng/µL) at 68°C overnight. Antidigoxigenin AP-conjugated antibody (1:5000, Roche 11093274910) was used to detect digoxigenin and was further stained with nitro-blue-tetrazolium/5-bromo-4-chloro-3-indolyl-phosphate (NBT/BCIP) solution (Roche, 11681451001).

Microinjection

Zebrafish embryos were microinjected with 8 ng *cbs* morpholino oligonucleotide (MO), 0.5 ng *cth* MO, or equivalent Ctrl MO at one-cell stage. The MOs were purchased from Gene Tools (Philomath, Oregon, USA), and the sequences are as follows.

cbs MO: 5'-CTGGCATGGTTTACCCTGACTATCA-3'
cth MO: 5'-GGCTGAGCTGTCGTTCTGCATCTCT-3'
 Ctrl MO: 5'-CCTCTTACCTCAGTTACAATTATA-3'

CRISPR/Cas9-mediated mutation of zebrafish *cbs* or *cth*

Mutations of zebrafish *cbs* or *cth* genes were induced by using the clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9 system as previously reported.^{22–24} Single guide RNA (sgRNA) sequences (*cbs*: 5'-GATGCTGACGATGTCTTCAA-3', *cth*: 5'-CGTTG-GTTCAGAGCCCCGAGC-3') were designed to target the sequences of mature *cbs* and *cth*. The paired oligonucleotides which contained the *cbs* or *cth* sgRNA sequence were annealed and cloned into a pT7-gRNA plasmid. Then, *cbs* sgRNA and *cth* sgRNA were synthesised with the MAXIscript T7 kit (Ambion, AM1312M). The synthesised sgRNA was purified with the mirVana miRNA isolation kit (Ambion, AM1560).²⁵ Coinjection of 600 pg Cas9 nuclease (NEB, M0386M) and 80 pg *cbs* or *cth* gRNA into zebrafish embryos were conducted at one-cell stage. The *cbs* or *cth* gene mutations in F0 embryos were examined by sequencing analysis after in vivo confocal imaging of the brain vasculature. The imaging data of F0 embryos carrying *cbs* or *cth* mutations were analysed.

The *cbs* or *cth* heterozygous mutant lines were generated by Nanjing XinJia Medical Technology Co. The two lines were separately outcrossed with the Tg(Flk1:eGFP) line. The crossed lines were further incrossed to observe the vascular development in homozygous mutant larvae at 3 and 5 dpf larvae. Genotyping was conducted after in vivo confocal imaging of the brain vasculature. The imaging data of embryos carrying *cbs* or *cth* homozygous mutations were analysed.

Drug treatment

GGY4137 (SIGMA, SML0100) was dissolved in zebrafish culture medium at 1 dpf (2000 µM for *cbs* MO group and 1000 µM for *cth* MO group). The medium was refreshed once a day and phenotypes were characterised at 3–5 dpf.

Measurement of NO production

Total NO production in the zebrafish larvae was examined with the total NO assay kit (Beyotime Institute of Biotechnology, S0023). It measured the concentration of nitrate and nitrite which are stable metabolites of NO.

Data analysis

All data were represented as mean±SEM. Statistical analysis was performed by using unpaired two-tailed Student's t-test or One-way analysis of variance.

RESULTS

Endogenous H₂S is necessary for the brain vascular development in zebrafish

To examine the expression pattern of *cbs* and *cth* in zebrafish, we conducted whole-mount in situ hybridisation in larval zebrafish at 3 dpf and found that both *cbs* and *cth* were ubiquitously expressed in the zebrafish brain (figure 1A). To examine whether H₂S is involved in brain vascular development, we first downregulated the expression of *cbs* and *cth* using MO-based knock-down. The expression of *cbs* and *cth* in zebrafish larvae were efficiently downregulated by the MOs (online supplemental figure S1) and H₂S production were also significantly reduced in *cbs* and *cth* morphants (online supplemental figure S2). We performed in vivo long-term serial confocal imaging of the midbrain vasculature during 3–5 dpf in the same transgenic zebrafish Tg(Flk1:eGFP)²⁶ larva, in which vascular ECs express enhanced green fluorescent protein (eGFP). Compared with embryos injected with a control MO ('Ctrl MO'), both the *cbs* and *cth* morphants displayed impaired brain vascular development (figure 1B, online supplemental figure S3A), as evidenced by reduced midbrain vessel density during 3–5 dpf (figure 1C; p<0.01). Meanwhile, the brain size (indicated by the width of the optic tectum²⁷) and cell apoptosis in *cbs* and *cth* morphants showed no significant difference compared with the Ctrl group at 3 dpf (online supplemental figure S4A,B,D and F), indicating that the impaired brain vascular development in *cbs* and *cth* morphants were not caused by the delayed brain development or





Figure 1 Knockdown and knockout of *cbs* or *cth* impair brain vascular development of larval zebrafish. (A) In situ hybridisation of whole zebrafish larvae at 3 dpf showing the ubiquitous expression of *cbs* and *cth* in the brain (lateral view). (B and C) Effects of morpholino oligonucleotide-mediated *cbs* or *cth* knockdown on brain vascular development. (B) Representative midbrain vessel structures reconstructed from confocal images of *cbs* or *cth* morphants at 3–5 dpf of the same larva. (C) Summary of data. The experiments were repeated three times, and six embryos were examined for each group at each time. (D and E) Effects of *cbs* or *cth* knockout (F0) on brain vascular development. (D) Representative midbrain vessel structures reconstructed from confocal images of the same F0 mutant larva at 3–5 dpf. (E) Summary of data. Six embryos were examined for each group. (F and G) Effects of *cbs* or *cth* homomutants on brain vascular development. (F) Representative midbrain vessel structures reconstructed from confocal images of the same *cbs* or *cth* homomutant larva at 3 and 5 dpf. (G) Summary of data. Six embryos were examined for each group. Scale bar, 300 μ m (A), 25 μ m (B, D, F). Error bars, SEM. * $P < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (unpaired two-tailed Student's *t*-test).

neuron death. Moreover, *cbs* and *cth* mutations (online supplemental figure S5) created by coinjecting *cbs* or *cth* sgRNA and Cas9 nuclease also caused a significant decrease in the midbrain vessel density during 3–5 dpf in F0 embryos (figure 1D,E, online supplemental figure S3B; $p < 0.01$). Furthermore, *cbs* and *cth* homozygous mutants (online supplemental figure S6), obtained by incrossing the stable heterozygous mutant lines with the Tg(Flk1:eGFP) background, also displayed impaired brain vascular development at 3 dpf, with the relative normal vascular development at 5 dpf possibly due to the compensatory effect of the stable lines (figure 1F,G, online supplemental figure S3C; $p < 0.05$). Although brains of *cbs* and *cth* homomutants were slightly smaller


at 3 dpf online supplemental figure S4C; $p < 0.05$), they did not show increased cell apoptosis (online supplemental figure S4E,G).

Next, we conducted rescue experiments on *cbs* and *cth* morphants by bath application of GYY4137 (200 μ M), a slow-releasing H_2S donor which is able to yield stable levels of H_2S at about 5–10 μ M at least for 7 days.²⁸ Embryos injected with MOs were exposed to Hanks' solution diluted with GYY4137 since 24 hours post fertilisation. GYY4137 treatment alleviated the brain vascular developmental defects in both *cbs* and *cth* morphants (figure 2). It is worth mentioning that administration of GYY4137 did not increase the midbrain vessel density in control fish (online supplemental figure S7). Taken

A



B



C




Figure 2 Rescue effect of GYY4137 on the defects of brain vascular development in *cbs* and *cth* morphants. (A) Representative projected confocal images showing that GYY4137 treatment ameliorated the impaired brain vascular development in *cbs* and *cth* morphants. Confocal images were taken at 3–5 dpf of the same larva. (B) Summary of the rescue effect of GYY4137. Six embryos were examined for each group. Scale bar, 50 μ m (A). Error bars, SEM. * $P < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (one-way analysis of variance). MO, morpholino oligonucleotide.

together, these results indicate that endogenous H_2S is important for the brain vascular development.

H_2S promotes brain angiogenesis rather than topological structure formation

Maturation of vascular networks involves complicated processes including vasculogenesis, angiogenesis and followed with vessel formation, stabilisation, branching, pruning and specialisation.²⁹ It can be generalised into two major aspects: vessel growth (via angiogenesis) and topological structure formation (ie, branching and pruning). To figure out the detailed roles of H_2S , we thus analysed quantitatively the characteristics of the midbrain vasculature. We used the total length and vessel segment number to, respectively, quantify vessel elongation and new vessel addition through angiogenesis, and the average order of vessel segments and vessel loop number to quantify the complexity of the vascular network.¹⁹ Both the total length and vessel segment number were markedly decreased in 3-dpf *cbs* and *cth* morphants (figure 3A–C; $p < 0.001$), while the average order and loop number had no significant change (figure 3A,D and E). These results indicate that H_2S enhances angiogenesis without affecting the topological structure during brain vascular development in zebrafish.

NOS/NO pathway is involved in the proangiogenic effect of H_2S

NO is formed from the reaction of L-arginine with O_2 , a reaction catalysed by NOS. Three NOS isoforms have been identified in mammals³⁰: neuronal NOS (NOS1), inducible NOS (NOS2) and endothelial NOS (NOS3). While in the zebrafish genome, there are one *nos1* homolog gene and two *nos2* homolog genes (*nos2a* and *nos2b*).³¹ We found that there was a decrease in *nos2a* expression in both *cbs* and *cth* morphants, but an increase in *nos2b* in *cth* morphants which might be due to a compensatory effect (figure 4A; $p < 0.05$). Meanwhile, the expression of *nos1* was not affected (figure 4A). Consistently, NO production was also decreased in both *cbs* and *cth* morphants (figure 4B; $p < 0.01$). We then performed rescue experiments by coinjection of *cbs* or *cth* MO with full-length zebrafish *nos2a* mRNA, and found that the decreased NO production and brain vascular defects caused by *cbs* or *cth* knockdown were significantly alleviated (figure 4C,D; $p < 0.01$ for NO production, $p < 0.05$ for vascular density). It is noteworthy that the expression of ERK1/2 (p44/42 mitogen-activated protein kinase), phosphorylated ERK1/2 and vascular endothelial growth factor (VEGF) was not significantly changed in both *cbs* and *cth* morphants (figure 4E,F). Taken together, these results suggest that the NOS/NO signalling pathway plays an important role in the proangiogenic effect of H_2S in brain vascular development.

DISCUSSION

H_2S has been shown to exert potent proangiogenic effects in vitro and in pathological models.^{14 15 32–35} Recently, two in vivo studies showed the importance of H_2S in muscle




Figure 3 Structure analysis of the midbrain vasculature in *cbs* and *cth* morphants. (A) Image of a 3-dpf larva showing the midbrain position delineated with dashed lines (upper left), and representative midbrain vasculature centerlines of 3-dpf larvae of Ctrl morpholino oligonucleotide (MO; upper right), *cbs* MO (lower left), and *cth* MO (lower right). C, caudal; L, lateral. (B–E) Summary of changes in the total vessel length (B), vessel segment number (C), weighted average segment Strahler order (D) and internal vessel loop number (E) of the midbrain vasculature in groups of Ctrl MO, *cbs* MO and *cth* MO. Six embryos were examined for each group. Error bars, SEM. *** $P < 0.001$ (unpaired two-tailed Student's t-test).

angiogenesis.^{16 17} By in vivo confocal imaging of zebrafish larvae, our study reveals the role of endogenous H_2S in developmental angiogenesis of the brain.

The signalling pathways involved in the proangiogenic effect of exogenous H_2S on peripheral vessels has been well studied with in vitro preparations.^{14 15 32–34} The proangiogenic effect of H_2S is mediated via: (1) EC-related angiogenic properties through a K_{ATP} channel/MAPK




Figure 4 The nitric oxide synthase (NOS)/nitric oxide (NO) pathway is involved in *cbs* and *cth* knockdown-induced defects of brain vascular development. (A and B) Effects of *cbs* and *cth* knockdown on the NOS/NO signalling pathway. Summary of *nos1*, *nos2a* and *nos2b* RNA expression (A) and total NO production (B) in 3-days post fertilisation (dpf) embryos. (C and D) Rescue effect of *nos2a* mRNA on the brain vascular developmental defects in *cbs* and *cth* morphants. Summary of increased total NO production (C) and midbrain vessel density (D) in 3-dpf embryos coinjected with *cbs* or *cth* MO and *nos2a* mRNA in comparison with those injected with *cbs* or *cth* MO. Six embryos were examined for each group. (E and F) No effect of *cbs* and *cth* knockdown on the protein expression of p-ERK1/2, ERK1/2 and VEGF. Representative blots (E) and summary (F) of Western blotting data. Error bars, SEM. * $P < 0.05$, ** $p < 0.01$ (unpaired two-tailed Student's t-test).

pathway¹⁴; (2) upregulation of VEGF expression and release from cells^{36 37}; (3) stimulation of the NOS/NO signalling pathway. In the present study, we examined ERK1/2 phosphorylation and VEGF expression and found no change of these signals in *chs* and *cth* morphants. NO and H₂S share many similar regulatory roles including promotion of angiogenesis, vasodilation, attenuation of apoptosis and antioxidant actions.^{32 38} H₂S is reported to exert proangiogenic functions via both NO-dependent and NO-independent mechanisms.^{32 34} Exogenously administered H₂S increases NOS expression³³ and phosphorylation³⁴ in cultured ECs and stimulates the production of NO. In zebrafish larvae, expression of *nos2a* was mainly detected in the head, eyes and gut regions,³¹ consistent with the expression pattern of *chs* and *cth* we observed. In the present study, depletion of H₂S resulted in a reduction in *nos2a* expression and NO production. Injection of full-length zebrafish *nos2a* mRNA alleviated the brain vascular developmental defects in *chs* and *cth* morphants. Thus, H₂S promotes brain angiogenesis via the NOS/NO pathway in zebrafish.

Studies demonstrating the proangiogenic effect of endogenous H₂S were mostly conducted through manipulating CTH,^{14 17 39} and the cellular target of H₂S is believed to be ECs as stated above. However, our study found that CBS-derived and CTH-derived H₂S both played vital roles in brain developmental angiogenesis. Indeed, in the brain, H₂S production in the astrocyte is almost 10 times as much as that in other brain cell types,⁶ making the astrocyte an important target for H₂S functioning and CBS a significant source of H₂S in the brain. We speculate that H₂S promotes brain angiogenesis by functioning on ECs through direct and indirect ways. The cellular mechanisms of H₂S promoting brain angiogenesis require further study.

Currently, numerous animal and cellular studies have showed that H₂S impacts ischaemic stroke outcomes due to its neuroprotective abilities.⁴⁰ Taking advantage of slow-releasing H₂S donors, H₂S is emerging as a potential therapy for cerebral ischaemia. Our study reveals the role of endogenous H₂S in brain angiogenesis, making H₂S-based therapies for cerebral ischaemia even more promising.

Contributors YW contributed to study design and interpretation of experimental data. WJ contributed to experiment performance and manuscript drafting. CL and MD contributed to analysis of experimental data. FW and XR contributed to manuscript revision. YF contributed to interpretation of experimental data. JD contributed to study design and final revision of the manuscript.

Funding The authors are indebted to Bing Xu and Yu Zhang for help with research design and technical supports. This work was supported by a grant from the Natural Science Foundation of China (31770800, 91849104, 81870274, 81571329).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval All animal use and handling procedures were approved by Institute of Neuroscience, Chinese Academy of Sciences.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. No additional unpublished data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Weiqing Jiang <http://orcid.org/0000-0003-1895-9618>

REFERENCES

- Goodwin LR, Francom D, Dieken FP, *et al.* Determination of sulfide in brain tissue by gas dialysis/ion chromatography: postmortem studies and two case reports. *J Anal Toxicol* 1989;13:105–9.
- Wang R. Two's company, three's a crowd: can H₂S be the third endogenous gaseous transmitter? *FASEB J* 2002;16:1792–8.
- Kimura H. Hydrogen sulfide: from brain to gut. *Antioxid Redox Signal* 2010;12:1111–23.
- Abe K, Kimura H. The possible role of hydrogen sulfide as an endogenous neuromodulator. *J Neurosci* 1996;16:1066–79.
- Robert K, Vialard F, Thiery E, *et al.* Expression of the cystathionine beta synthase (CBS) gene during mouse development and immunolocalization in adult brain. *J Histochem Cytochem* 2003;51:363–71.
- Lee M, Schwab C, Yu S, *et al.* Astrocytes produce the antiinflammatory and neuroprotective agent hydrogen sulfide. *Neurobiol Aging* 2009;30:1523–34.
- Enokido Y, Suzuki E, Iwasawa K, *et al.* Cystathionine beta-synthase, a key enzyme for homocysteine metabolism, is preferentially expressed in the radial glia/astrocyte lineage of developing mouse CNS. *FASEB J* 2005;19:1854–6.
- Kimura H. Hydrogen sulfide: its production, release and functions. *Amino Acids* 2011;41:113–21.
- Leffler CW, Parfenova H, Basuroy S, *et al.* Hydrogen sulfide and cerebral microvascular tone in newborn pigs. *Am J Physiol Heart Circ Physiol* 2011;300:H440–7.
- Kabil O, Vitvitsky V, Xie P, *et al.* The quantitative significance of the transsulfuration enzymes for H₂S production in murine tissues. *Antioxid Redox Signal* 2011;15:363–72.
- Hu L-F, Lu M, Hon Wong PT, *et al.* Hydrogen sulfide: neurophysiology and neuropathology. *Antioxid Redox Signal* 2011;15:405–19.
- Yang G, Wu L, Jiang B, *et al.* H₂S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 2008;322:587–90.
- Cai W-J, Wang M-J, Moore PK, *et al.* The novel proangiogenic effect of hydrogen sulfide is dependent on Akt phosphorylation. *Cardiovasc Res* 2007;76:29–40.
- Papapetropoulos A, Pyriochou A, Altaany Z, *et al.* Hydrogen sulfide is an endogenous stimulator of angiogenesis. *Proc Natl Acad Sci U S A* 2009;106:21972–7.
- Wang M-J, Cai W-J, Li N, *et al.* The hydrogen sulfide donor NaHS promotes angiogenesis in a rat model of hind limb ischemia. *Antioxid Redox Signal* 2010;12:1065–77.
- Das A, Huang GX, Bonkowski MS, *et al.* Impairment of an Endothelial NAD⁺-H₂S Signaling Network Is a Reversible Cause of Vascular Aging. *Cell* 2018;173:74–89.
- Longchamp A, Mirabella T, Arduini A, *et al.* Amino Acid Restriction Triggers Angiogenesis via GCN2/ATF4 Regulation of VEGF and H₂S Production. *Cell* 2018;173:117–29.
- Weinstein B. Vascular cell biology in vivo: a new piscine paradigm? *Trends Cell Biol* 2002;12:439–45.
- Chen Q, Jiang L, Li C, *et al.* Haemodynamics-driven developmental pruning of brain vasculature in zebrafish. *PLoS Biol* 2012;10:e1001374.

- 20 Wienholds E, Kloosterman WP, Miska E, *et al.* MicroRNA expression in zebrafish embryonic development. *Science* 2005;309:310–1.
- 21 Yu P-chun, Gu S-ye, Bu J-wen, *et al.* Trpc1 is essential for in vivo angiogenesis in zebrafish. *Circ Res* 2010;106:1221–32.
- 22 Li J, Zhang B-bing, Ren Y-gang, *et al.* Intron targeting-mediated and endogenous gene integrity-maintaining knockin in zebrafish using the CRISPR/Cas9 system. *Cell Res* 2015;25:634–7.
- 23 Rossi A, Kontarakis Z, Gerri C, *et al.* Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature* 2015;524:230–3.
- 24 Shah AN, Davey CF, Whitebitch AC, *et al.* Rapid reverse genetic screening using CRISPR in zebrafish. *Nat Methods* 2015;12:535–40.
- 25 Xu B, Zhang Y, Du X-F, *et al.* Neurons secrete miR-132-containing exosomes to regulate brain vascular integrity. *Cell Res* 2017;27:882–97.
- 26 Roman BL, Pham VN, Lawson ND, *et al.* Disruption of ACVRL1 increases endothelial cell number in zebrafish cranial vessels. *Development* 2002;129:3009–19.
- 27 Näslund J. A simple non-invasive method for measuring gross brain size in small live fish with semi-transparent heads. *PeerJ* 2014;2:e586.
- 28 Lee ZW, Zhou J, Chen C-S, *et al.* The slow-releasing hydrogen sulfide donor, GYY4137, exhibits novel anti-cancer effects in vitro and in vivo. *PLoS One* 2011;6:e21077.
- 29 Lee HS, Han J, Bai H-J, *et al.* Brain angiogenesis in developmental and pathological processes: regulation, molecular and cellular communication at the neurovascular interface. *Febs J* 2009;276:4622–35.
- 30 Knowles RG, Moncada S. Nitric oxide synthases in mammals. *Biochem J* 1994;298:249–58.
- 31 Lepiller S, Franche N, Solary E, *et al.* Comparative analysis of zebrafish nos2a and nos2b genes. *Gene* 2009;445:58–65.
- 32 Holwerda KM, Karumanchi SA, Lely AT. Hydrogen sulfide: role in vascular physiology and pathology. *Curr Opin Nephrol Hypertens* 2015;24:170–6.
- 33 Bir SC, Kolluru GK, McCarthy P, *et al.* Hydrogen sulfide stimulates ischemic vascular remodeling through nitric oxide synthase and nitrite reduction activity regulating hypoxia-inducible factor-1 α and vascular endothelial growth factor-dependent angiogenesis. *J Am Heart Assoc* 2012;1:e004093.
- 34 Coletta C, Papapetropoulos A, Erdelyi K, *et al.* Hydrogen sulfide and nitric oxide are mutually dependent in the regulation of angiogenesis and endothelium-dependent vasorelaxation. *Proc Natl Acad Sci U S A* 2012;109:9161–6.
- 35 Madurga A, Golec A, Pozarska A, *et al.* The H₂S-generating enzymes cystathionine β -synthase and cystathionine γ -lyase play a role in vascular development during normal lung alveolarization. *Am J Physiol Lung Cell Mol Physiol* 2015;309:L710–24.
- 36 Kan J, Guo W, Huang C, *et al.* S-Propargyl-Cysteine, a novel water-soluble modulator of endogenous hydrogen sulfide, promotes angiogenesis through activation of signal transducer and activator of transcription 3. *Antioxid Redox Signal* 2014;20:2303–16.
- 37 Holwerda KM, Burke SD, Faas MM, *et al.* Hydrogen sulfide attenuates sFlt1-induced hypertension and renal damage by upregulating vascular endothelial growth factor. *J Am Soc Nephrol* 2014;25:717–25.
- 38 Polhemus DJ, Lefer DJ. Emergence of hydrogen sulfide as an endogenous gaseous signaling molecule in cardiovascular disease. *Circ Res* 2014;114:730–7.
- 39 Tao B-B, Liu S-Y, Zhang C-C, *et al.* Vegfr2 functions as an H₂S-targeting receptor protein kinase with its novel Cys1045-Cys1024 disulfide bond serving as a specific molecular switch for hydrogen sulfide actions in vascular endothelial cells. *Antioxid Redox Signal* 2013;19:448–64.
- 40 Jia J, Li J, Cheng J. H₂S-based therapies for ischaemic stroke: opportunities and challenges. *Stroke Vasc Neurol* 2019;4:63–6.