

Association of APOE genotype with CT markers of cerebral amyloid angiopathy in spontaneous intracerebral haemorrhage

Qiong Yang,¹ Xiangzhu Zeng,² Lu Tang,¹ Xiaolu Liu,¹ Kailin Xia,¹ Feng Gao,³ Xu Huang,³ Nan Li,⁴ Dongsheng Fan ^{1,5,6}

ABSTRACT

To cite: Yang Q, Zeng X, Tang L, et al. Association of APOE genotype with CT markers of cerebral amyloid angiopathy in spontaneous intracerebral haemorrhage. Stroke & Vascular Neurology 2024;0. doi:10.1136/ svn-2024-003477

Additional supplemental material is published online only. To view, please visit the journal online (https://doi.org/10.1136/ svn-2024-003477).

QY and XZ contributed equally.

QY and XZ are joint first authors.

Received 12 June 2024 Accepted 21 November 2024

C Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Check for updates

For numbered affiliations see end of article.

Correspondence to

Dongsheng Fan; dsfan@sina.com

Background and objective We investigated the association of APOE alleles with CT-based cerebral amyloid angiopathy (CAA) markers including subarachnoid extension (SAE) and finger-like projection (FLP). Methods We included patients with acute primary supratentorial intracerebral haemorrhage (ICH) from a multicentre cohort in China. First, the association of APOE with ICH location (lobar vs non-lobar) was evaluated. Next, the relationships of APOE with SAE, FLP, and the coexistence of the two (SAE+FLP) were evaluated. Results 533 patients with supratentorial ICH were enrolled. Among them were 138 patients with lobar ICH and 395 with non-lobar ICH. Compared with the non-lobar group, APOE £4 (OR 1.894, 95% CI 1.138 to 3.154, p=0.014) and ɛ2/ɛ4 (OR 6.098, 95% Cl 1.414 to 26.293, p=0.015) were associated with lobar ICH. With regard to CAA markers, APOE E2 was associated with SAE (OR 2.109, 95% CI 1.167 to 3.810, p=0.013), ɛ4 was associated with FLP and SAE+FLP (OR 3.026, 95% CI 1.353 to 6.767, p=0.007; OR 3.514, 95% CI 1.485 to 8.316, p=0.004, respectively) and $\varepsilon 2/\varepsilon 4$ was associated with all three factors (SAH: OR 7.599, 95% CI 1.764 to 32.734, p=0.006; FLP: OR 20.333, 95% CI 3.278 to 126.137, p=0.001; SAE+FLP: OR 30.568, 95% CI 4.460 to 209.503, p<0.001) after adjusting for age, and remained significant after adjusting for age and ICH volume. **Conclusion** In patients with spontaneous supratentorial ICH, APOE ε 2 and ε 4 alleles were associated with SAE and FLP, respectively, suggesting APOE allele-specific effects on CT markers of CAA and their potential mechanisms.

BACKGROUND Spontaneous

intracerebral haemorrhage (ICH) is a critical disease which accounts for 20% of all strokes, with high mortality and morbidity (approximately 50% and 30%, respectively).¹² Incidence of ICH is expected to remain significant with the ageing of the population and the extensive use of anticoagulant drugs.³

The majority of primary ICH are caused by two common cerebral small vessel diseases, arteriolosclerosis and cerebral amyloid angiopathy (CAA).³ Arteriolosclerosis, related to

WHAT IS ALREADY KNOWN ON THIS TOPIC

 \Rightarrow Subarachnoid extension (SAE) and finger-like proiection (FLP) may be biomarkers for the risk and severity of cerebral amyloid angiopathy (CAA)associated intracerebral haemorrhage (ICH), but the mechanism of these neuroimaging manifestations is still unclear, and clarifying the associations between apolipoprotein E (APOE) genotype and SAE or FLP may be helpful.

WHAT THIS STUDY ADDS

 \Rightarrow In patients with supratentorial ICH, APOE ε 2 was associated with SAE, APOE ε 4 was associated with FLP and SAE+FLP and $\epsilon 2/\epsilon 4$ heterozygosity was associated with all three.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

 \Rightarrow This study revealed that different APOE alleles have different effects on CT markers of CAA, suggesting different pathophysiological processes.

hypertension, often involves the perforator arteries of the basal ganglia and brainstem and thus causes deep territories bleeding; CAA mainly refers to the deposition of β-amyloid in the walls of arterioles and capillaries in the pia mater and cortex, which causes lobar ICH.³ The risk of recurrence in patients with CAA-associated ICH is 7.4%/year, which is greater than that of non-CAA-associated ICH patients (1.1%/year).³

CAA imaging markers based on CT have been developed.⁴ The pathologically validated Edinburgh CT diagnostic criteria include the subarachnoid extension (SAE) and fingerlike projection (FLP).5-7 In patients with CAA, SAE seems to be associated with the increased risk of incidence and recurrence of ICH.⁸⁻¹¹ Compared with patients with SAE or FLP alone, patients with SAE combined with FLP have more severe ICH at onset, higher in-hospital case-fatality and higher mRS scores at discharge.¹²Thus, SAE and FLP may



be biomarkers for the risk and severity of CAA-associated ICH, but the mechanism of these neuroimaging manifestations is still unclear, and clarifying the associations between apolipoprotein E (APOE) genotype and SAE or FLP may be helpful. The APOE genotype is an important genetic risk factor for ICH and associated with the prognosis.^{13–15} The APOE ɛ4 allele is associated with CAA, while APOE ε_2 is associated with the severity of ICH.^{13 16 17} The two APOE alleles may play different roles in the pathogenesis of CAA, that is, APOE ε 4 promotes the deposition of β -amyloid in blood vessels, APOE ϵ 2 increases the damage of vessel wall caused by amyloid deposition and the $\epsilon 2/\epsilon 4$ genotype may represent a double hit to superficial blood vessels, which promotes both deposition of amyloids and also wall rupture of the most fragile arterioles in CAA.^{18 19} APOE is speculated to also mediate the selective distribution of β -amyloid, that is, APOE ϵ 4 favours cortical CAA and £2 is associated with leptomeningeal CAA.²⁰ Cortical superficial siderosis (cSS, considered as a chronic manifestation of SAE) is associated with $APOE \varepsilon 2$, ^{18 20–22} and an association between SAE and ɛ2 has also been observed in patients with supratentorial ICH, though these data were obtained from small studies.²³ As for the relationship between FLP and APOE, only one study has shown that FLP in patients with lobar ICH is associated with $\varepsilon 4$ and $\epsilon 2/\epsilon 4$ heterozygosity; surprisingly, no association was observed between SAE and APOE in this study.²⁴

To date, there is no consistent conclusion about the association between *APOE* and SAE or FLP. On this basis, we hypothesised that different *APOE* alleles were associated with these two markers, which we tested in an ICH patient cohort.

METHODS

Research settings

We included patients from a prospective multicentre cohort of acute primary ICH patients recruited from 19 hospitals in Beijing, Hebei and Inner Mongolia between 2015 and 2019 (NCT number (REDACTED)). The main inclusion criteria of this study were as follows: (1) primary spontaneous supratentorial ICH and (2) available *APOE* genotype. Patients were excluded if they had any of the following characteristics: (1) secondary haemorrhage, such as vascular malformation, trauma, tumour or haemorrhagic transformation in ischaemic stroke; (2) absence of non-contrast CT within 7 days after the onset of symptoms or low-quality image or (3) head surgery performed before baseline CT.

Data collection

The demographic and clinical data were systematically and prospectively collected and recorded by trained neurologist at the time of index symptomatic ICH and included age, sex, vascular risk factors, patient medication history before ICH, previous ICH history and family history according to the medical records or description from patients or their relatives. The National Institutes of Health Stroke Scale (NIHSS) Score at admission was evaluated.

Genotyping

The *APOE* genotype was determined from blood samples donated by patients at enrolment. DNA was extracted from blood samples. Two loci of the *APOE* gene (rs429358 and rs7412) were detected and translated into *APOE* genotypes ($\varepsilon_3/\varepsilon_3, \varepsilon_3/\varepsilon_4, \varepsilon_4/\varepsilon_4, \varepsilon_3/\varepsilon_2, \varepsilon_2/\varepsilon_2$ and $\varepsilon_2/\varepsilon_4$). Genotyping was performed by laboratory staff blinded to the clinical and neuroimaging data. *APOE* alleles were classified by their presence or absence (dominant model) and allele count (additive model) and $\varepsilon_2/\varepsilon_4$ heterozygosity was observed.²⁴

Image acquisition and analysis

CT scans were reviewed by radiologists blinded to both the clinical data and *APOE* genotype to determine the location and volume of ICH and the presence or absence of intraventricular haemorrhage. The location of ICH was defined as supratentorial (lobar and deep), or infratentorial (brainstem and cerebellum) using the cerebral haemorrhage anatomical rating instrument (CHARTS).²⁵ The volume of ICH was calculated from the baseline CT images using the ABC/2 method.

The presence of SAE and FLP was assessed in the CT axial plane according to the Edinburgh diagnostic criteria and both written and pictorial definitions were used to improve accuracy; SAE was defined as an extra-axial haemorrhage in the subarachnoid space, while FLP was defined as elongated extensions arising from the haematoma, longer than wide, with or without the extension to the cortex (figure 1).⁵ The assessment was independently performed by two experienced investigators (QY and XZ) also blinded to both the clinical data and *APOE* genotype. The inter-rater agreement was good for cSAH presence (kappa=0.756), as well as for FLP (kappa=0.780). Discrepancies were settled by consensus after the investigators read all of the scans. Patients with CT within 7 days after the onset of symptoms were included, as patients with



Figure 1 Examples of subarachnoid extension (SAE, black arrow) and finger-like projections (FLP, white arrow). FLP, finger-like projection; SAE, subarachnoid extension.

CT up to 7 days were included in the study of Edinburgh criteria.⁵

Statistical analysis

Due to the nature of the underlying distributions, continuous variables are presented as medians with IQR (first quartile to third quartile). Categorical variables are expressed as n (%). First, univariate and age-adjusted multivariate logistic regression analyses were used to evaluate the association between APOE and ICH location (non-lobar vs lobar). Second, in all supratentorial ICH patients, the association between APOE and the presence of CAA neuroimaging markers (ie, SAE, FLP and SAE+FLP), using participants without either SAE or FLP as the comparator group, was analysed using univariate and multivariate logistic regressions (adjusting for age, and adjusting for age and ICH volume, respectively). Third, factors associated with SAE, FLP, SAE+FLP were identified using multivariable analysis including two kinds of models. In model 1, prespecified plausible predictors (including age, ICH volume, ICH location and APOE) as well as variables with a p value<0.1 in univariable regression were included for multivariable logistic regression. In model 2, same variables as model 1 were included, and backward (conditional) logistic regression was used. Fourth, subgroup analyses by location that is, lobar ICH and non-lobar ICH were further performed. Statistical significance was defined as p<0.05. All analyses were performed with SPSS (V.26.0).

RESULTS

Study population

A total of 533 patients were eligible for analysis, with a median age of 61 (52, 74) years and 355 males (66.6%). Among them were 138 patients with lobar ICH and 395 patients with non-lobar ICH. Figure 2 shows the flow



Figure 2 Study flowchart. ; APOE, apolipoprotein E; ICH intracerebral haemorrhage.

location, APOE ε 4 as a dominant variable was associated with FLP (OR 2.886, 95% CI 1.011 to 8.232, p = 0.048), while neither APOE $\varepsilon 4$ as an additive variable nor APOE $\epsilon 2/\epsilon 4$ was associated with FLP significantly (overall p =

chart of patient inclusion. Compared with the included patients, the excluded patients had more hypertension, and fewer family history of ICH, while there were no significant differences in remaining characteristics (online supplemental table S1).

The baseline characteristics and the APOE allele frequency of CT markers including SAE and FLP of all patients and patients by ICH location are shown in tables 1 and 2. SAE, FLP and SAE+FLP were present in 86 (16.1%), 37 (6.9%) and 31 (5.8%) patients, respectively. Compared with the non-lobar ICH group, the lobar ICH group were older, had fewer hypertension, more previous ICH and greater ICH volume, as well as more SAE, FLP and SAE+FLP (table 1). The characteristics of patients with or without SAE, FLP and SAE+FLP were shown in online supplemental table S2.

APOE and ICH location

Compared with non-lobar ICH, APOE E4 as a dominant variable was associated with lobar ICH (OR 1.894, 95% CI 1.138 to 3.154, p=0.014) in the age-adjusted multivariate analysis. This risk increased with the increasing allele count (additive model, overall p=0.004). APOE $\varepsilon 2/\varepsilon 4$ was also associated with lobar ICH (OR 6.098, 95% CI 1.414 to 26.293, p=0.015) (table 3).

APOE and imaging markers in all ICH APOE and SAE in all ICH

The APOE genotype associated with SAE included APOE ε2 as a dominant variable (OR 2.109, 95% CI 1.167-3.810, p = 0.013), an additive variable (overall p = 0.046) and APOE $\varepsilon 2/\varepsilon 4$ (OR 7.599, 95% CI 1.764-32.734, p = 0.006) after adjusting for age, and remained significant after adjusting for age and ICH volume (table 4).

In a sensitivity analysis that adjusting for age, hypertension, smoker, previous ICH, ICH volume, ICH location and intraventricular haemorrhage, APOE $\varepsilon 2$ as a dominant variable was associated with SAE (OR 2.436, 95% CI 1.094 to 5.425, p = 0.029), while neither APOE ε 2 as an additive variable nor APOE $\varepsilon 2/\varepsilon 4$ was associated with SAE significantly (overall p = 0.068; p = 0.197, respectively) in all supratentorial ICH patients (online supplemental table S3).

APOE and FLP in all ICH

The APOE genotype associated with FLP included APOE ε4 as a dominant variable (OR 3.026, 95% CI 1.353 to 6.767, p=0.007), and APOE $\varepsilon 2/\varepsilon 4$ (OR 20.333, 95% CI 3.278 to 126.137, p=0.001) after adjusting for age, and remained significant after adjusting for age and ICH volume (table 4).

In a sensitivity analysis that adjusting for age, hypertension, previous OAC use, previous ICH, ICH volume and

Baseline characteristics of patients with ICH (all, lobar and non-lobar)							
All supratentorial ICH (n=533)	Lobar ICH (n=138)	Non-lobar ICH (n=395)					
61 (52, 74)	71.5 (57, 80)	59 (50, 69)					
355/533 (66.6)	85/138 (61.6)	270/395 (68.4)					
360/521 (69.1)	72/138 (52.2)	288/383 (75.2)					
79/519 (15.2)	23/135 (17.0)	56/384 (14.6)					
129/509 (25.3)	31/129 (24.0)	98/380 (25.8)					
52/520 (10.0)	9/132 (6.8)	43/388 (11.1)					
2/512 (0.4)	1/131 (0.8)	1/381 (0.3)					
57/493 (11.6)	16/126 (12.7)	41/367 (11.2)					
34/497 (6.8)	10/123 (8.1)	24/374 (6.4)					
25/470 (5.3)	9/118 (7.6)	16/352 (4.5)					
68/518 (13.1)	29/136 (21.3)	39/382 (10.2)					
7 (2, 13)	3 (1, 12)	8 (3, 13)					
10.3 (4.4, 22.9)	20.2 (7.7, 45.2)	8 (4, 17.5)					
86/533 (16.1)	61/138 (44.2)	25/395 (6.3)					
37/533 (6.9)	30/138 (21.7)	7/395 (1.8)					
31/533 (5.8)	27/138 (19.6)	4/395 (1.0)					
168/524 (32.1)	39/138 (28.3)	129/386 (33.4)					
89/533 (16.7)	28/138 (20.3)	61/395 (15.4)					
85/533 (15.9)	26/138 (18.8)	59/395 (14.9)					
4/533 (0.8)	2/138 (1.4)	2/395 (0.5)					
512/533 (96.1)	124/138 (89.9)	388/395 (98.2)					
148/533 (27.8)	40/138 (29.0)	108/395 (27.3)					
364/533 (68.3)	84/138 (60.9)	280/395 (70.9)					
89/533 (16.7)	32/138 (23.2)	57/395 (14.4)					
81/533 (15.2)	26/138 (18.8)	55/395 (13.9)					
8/533 (1.5)	6/138 (4.3)	2/395 (0.5)					
9/533 (1.7)	6/138 (4.3)	3/395 (0.8)					
	All supratentorial ICH (n=533) 61 (52, 74) 355/533 (66.6) 360/521 (69.1) 79/519 (15.2) 129/509 (25.3) 52/520 (10.0) 2/512 (0.4) 57/493 (11.6) 34/497 (6.8) 25/470 (5.3) 68/518 (13.1) 7 7 (2, 13) 10.3 (4.4, 22.9) 86/533 (16.1) 37/533 (6.9) 31/533 (5.8) 168/524 (32.1) 89/533 (16.7) 85/533 (16.7) 85/533 (16.7) 85/533 (16.7) 84/533 (27.8) 364/533 (68.3) 89/533 (16.7) 81/533 (15.2) 8/533 (15.2) 8/533 (15.7) 8/533 (15.7) 8/533 (15.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (16.7) 8/533 (15.2) 8/533 (1.5) </td <td>All supratentorial ICH (n=533) Lobar ICH (n=138) 61 (52, 74) 71.5 (57, 80) 355/533 (66.6) 85/138 (61.6) 360/521 (69.1) 72/138 (52.2) 79/519 (15.2) 23/135 (17.0) 129/509 (25.3) 31/129 (24.0) 52/520 (10.0) 9/132 (6.8) 2/512 (0.4) 1/131 (0.8) 57/493 (11.6) 16/126 (12.7) 34/497 (6.8) 10/123 (8.1) 25/470 (5.3) 9/118 (7.6) 68/518 (13.1) 29/136 (21.3) 7 7(2, 13) 3 (1, 12) 10.3 (4.4, 22.9) 20.2 (7.7, 45.2) 86/533 (16.1) 61/138 (44.2) 37/533 (6.9) 30/138 (21.7) 31/533 (5.8) 27/138 (19.6) 168/524 (32.1) 39/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 84/533 (0.8) 4/533 (0.8) <!--</td--></td>	All supratentorial ICH (n=533) Lobar ICH (n=138) 61 (52, 74) 71.5 (57, 80) 355/533 (66.6) 85/138 (61.6) 360/521 (69.1) 72/138 (52.2) 79/519 (15.2) 23/135 (17.0) 129/509 (25.3) 31/129 (24.0) 52/520 (10.0) 9/132 (6.8) 2/512 (0.4) 1/131 (0.8) 57/493 (11.6) 16/126 (12.7) 34/497 (6.8) 10/123 (8.1) 25/470 (5.3) 9/118 (7.6) 68/518 (13.1) 29/136 (21.3) 7 7(2, 13) 3 (1, 12) 10.3 (4.4, 22.9) 20.2 (7.7, 45.2) 86/533 (16.1) 61/138 (44.2) 37/533 (6.9) 30/138 (21.7) 31/533 (5.8) 27/138 (19.6) 168/524 (32.1) 39/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 85/533 (16.7) 28/138 (20.3) 84/533 (0.8) 4/533 (0.8) </td					

*Moderate to severe alcohol consumption was defined as regular alcohol consumption ≥2 units per day (1 unit=360 mL of beer, 25 mL of 40% spirit or 100 mL of wine).

APOE, apolipoprotein E; FLP, finger-like projections; ICH, intracerebral haemorrhage; NIHSS, National Institutes of Health Stroke Scale; OAC, oral anticoagulation; SAE, subarachnoid extension.

0.109; p = 0.065, respectively) (online supplemental table S4).

APOE and SAE+FLP in all ICH

The *APOE* genotype associated with SAE+FLP included *APOE* ϵ 4 as a dominant variable (OR 3.514, 95% CI 1.485 to 8.316, p=0.004), *APOE* ϵ 4 as an additive variable (overall p=0.017) and *APOE* ϵ 2/ ϵ 4 (OR 30.568, 95% CI 4.460 to 209.503, p<0.001) in the age-adjusted multivariate analysis, and remained significant after adjusting for age and ICH volume (table 4).

In a sensitivity analysis that adjusting for age, hypertension, previous OAC use, previous ICH, ICH volume and location, APOE $\varepsilon 4$ as a dominant variable and APOE $\varepsilon 2/\varepsilon 4$, were associated with SAE+FLP (OR 3.612, 95% CI 1.110 to 11.748, p = 0.033; OR 13.005, 95% CI 1.290 to 131.118, p = 0.030, respectively)), while APOE $\varepsilon 2$ as an additive variable was not significantly associated with SAE+FLP (overall p = 0.075) (online supplemental table S5).

Table 2 AP	OE allele fre	quency of C	AA neuroima	aging biomarke	ers in patients	s with ICH (a	ll, lobar and	non-lobar)				
	All suprate	entorial ICH			Lobar ICH				Non-lobar I	сH		
	No SAE or FLP (441/533)	SAE (86/533)	FLP (37/533)	SAE+FLP* (31/533)	No SAE or FLP (74/138)	SAE (61/138)	FLP (30/138)	SAE+FLP* (27/138)	No SAE or FLP (367/395)	SAE (25/395)	FLP (7/395)	SAE+FI (4/395)
APOE ε2, N (%)											
Any allele	67 (15.2)	22 (25.6)	7 (18.9)	7 (22.6)	11 (14.9)	17 (27.9)	7 (23.3)	7 (25.9)	56 (15.3)	5 (20.0)	0 (0)	0 (0)
1 allele	64 (14.5)	21 (24.4)	7 (18.9)	7 (22.6)	10 (13.5)	16 (26.2)	7 (23.3)	7 (25.9)	54 (14.7)	5 (20.0)	0 (0)	0 (0)
2 alleles	3 (0.7)	1 (1.2)	(0) 0	(0) 0	1 (1.4)	1 (1.6)	0 (0)	(0) 0	2 (0.5)	0 (0)	0 (0)	0 (0)
APOE ε3, N(5	(%											
Any allele	427 (96.8)	79 (91.9)	33 (89.2)	27 (87.1)	67 (90.5)	54 (88.5)	26 (86.7)	23 (85.2)	360 (98.1)	25 (100.0)	7 (100.0)	4 (100.C
1 allele	118 (26.8)	29 (33.7)	12 (32.4)	11 (35.5)	17 (23.0)	22 (36.1)	11 (36.7)	10 (37.0)	101 (27.5)	7 (28.0)	1 (14.3)	1 (25.0)
2 alleles	309 (70.1)	50 (58.1)	21 (56.8)	16 (51.6)	50 (67.6)	32 (52.5)	15 (50.0)	13 (48.1)	259 (70.6)	18 (72.0)	6 (85.7)	3 (75.0)
APOE ε4, N((%)											
Any allele	69 (15.6)	19 (22.1)	12 (32.4)	11 (35.5)	14 (18.9)	17 (27.9)	11 (36.7)	10 (37.0)	55 (15.0)	2 (8.0)	1 (14.3)	1 (25.0)
1 allele	62 (14.1)	18 (20.9)	11 (29.7)	10 (32.3)	9 (12.2)	16 (26.2)	10 (33.3)	9 (33.3)	53 (14.4)	2 (8.0)	1 (14.3)	1 (25.0)
2 alleles	7 (1.6)	1 (1.2)	1 (2.7)	1 (3.2)	5 (6.8)	1 (1.6)	1 (3.3)	1 (3.7)	2 (0.5)	0 (0)	0 (0)	(0) 0
APOE ε2/ ε4, N(%)	4 (0.9)	5 (5.8)	3 (8.1)	3 (9.7)	1 (1.4)	5 (8.2)	3 (10.0)	3 (11.1)	3 (0.8)	(0) 0	0 (0)	(0) 0
*Participants ir CAA, cerebral	N SAE+FLP gr amyloid angic	roup were also opathy; ICH, ir	o included in th	ne separate SAE aemorrhage; AP0	and FLP grou <i>DE</i> , apolipopro	ps. otein E; SAE, s	ubarachnoid ∈	extension; FLP, fir	ngerlike projecti	ions.		

*

Open acce	SS					
Table 3 Asso	ociation of APOE w	ith ICH locatio	ns			
	Non-lobar ICH	Lobar ICH	Univariate		Multivariate (age adjuste	ed)
	(n=395)	(n=138)	OR (95% CI)	P value	OR (95% CI)	P value
APOE, N (%)						
APOE ε2 dom	inant					
Any allele	61 (15.4)	28 (20.3)	1.394 (0.848 to 2.290)	0.190	1.438 (0.855 to 2.417)	0.171
APOE ε2 addi	tive					
1 allele	59 (14.9)	26 (18.8)	1.338 (0.804 to 2.226)	0.306	1.384 (0.813 to 2.356)	0.303
2 alleles	2 (0.5)	2 (1.4)	3.036 (0.423 to 21.811)		3.062 (0.364 to 25.726)	
APOE ε4 dom	inant					
Any allele	57 (14.4)	32 (23.2)	1.790 (1.102 to 2.907)	0.019	1.894 (1.138 to 3.154)	0.014
APOE ε4 addi	tive					
1 allele	55 (13.9)	26 (18.8)	1.507 (0.901 to 2.523)	0.009	1.551 (0.903 to 2.664)	0.004
2 alleles	2 (0.5)	6 (4.3)	9.566 (1.902 to 48.102)		14.368 (2.557 to 80.717)	
APOE ε2/ε4	3 (0.8)	6 (4.3)	5.939 (1.465 to 24.082)	0.013	6.098 (1.414 to 26.293)	0.015

APOE and imaging markers in ICH subgroup by location

ICH, intracerebral haemorrhage; APOE, apolipoprotein E.

In patients with lobar ICH, the association between SAE and *APOE* ϵ 2 or ϵ 2/ ϵ 4 was not significant in the age and volume-adjusted multivariate analysis (ϵ 2 as a dominant variable: OR 2.704, 95% CI 0.865 to 8.454, p=0.087; ϵ 2/ ϵ 4: OR 9.333, 95% CI 0.739 to 117.837, p=0.084); *APOE* ϵ 4 as a dominant variable was associated with FLP (OR 3.083, 95% CI 1.164 to 8.164, p=0.023) after adjusting for age, while not significant after adjusting for age and ICH volume (OR 2.913, 95% CI 0.958 to 8.859, p=0.060); *APOE* ϵ 4 as a dominant variable was associated with SAE+FLP (OR 3.006, 95% CI 1.112 to 8.125, p=0.030) after adjusting for age, while not significant after adjusting for age and ICH volume (OR 2.951, 95% CI 0.915 to 9.517, p=0.070) (online supplemental table S6).

In patients with non-lobar ICH, there was no association observed between SAE, FLP or SAE+FLP and *APOE* in the univariate and multivariate analysis (online supplemental table S7).

DISCUSSION

This multicentre ICH cohort analysis found that in patients with supratentorial ICH, $APOE \epsilon 2/\epsilon 4$ heterozygosity and the $\epsilon 4$ allele were independently associated with lobar ICH compared with non-lobar ICH. The results of APOE and CT markers showed that $APOE \epsilon 2$ was associated with SAE, $APOE \epsilon 4$ was associated with FLP and SAE+FLP and $\epsilon 2/\epsilon 4$ heterozygosity was associated with all three. The results showed that different APOE alleles have different effects on CT markers of CAA, suggesting different pathophysiological processes.

APOE $\varepsilon4$ and $\varepsilon2/\varepsilon4$ were associated with lobar ICH compared with non-lobar ICH in this study, which is consistent with previous results and supports the *APOE* genotype as a strong genetic risk factor for CAA, which is prone to cause lobar ICH.²⁴

In this study, different CT neuroimaging markers exhibited different associations with the APOE alleles. The associations of SAE with APOE $\varepsilon 2$ and $\varepsilon 2/\varepsilon 4$ heterozygosity further validate and expand the results of previous studies.²³The associations of FLP with APOE ε 4 and ε 2/ ε 4 heterozygosity are consistent with a previous study, which mainly included Caucasians with lobar ICH.²⁴ Thus, our findings validate these results in Chinese patients with supratentorial ICH. FLP rarely occurs alone in patients with lobar ICH and is mostly accompanied by SAE.¹² Compared with patients with only one CT marker or none, patients with both SAE and FLP had more severe ICH (including higher ICH volume and higher NIHSS score) and a poorer prognosis (more in-hospital deaths and higher modified Rankin scale score).¹² When we further analysed the relationship between APOE and SAE+FLP, we found that $\varepsilon 4$ and $\varepsilon 2/\varepsilon 4$ heterozygosity were associated with it, which was not found in previous studies.

These findings suggest that APOE may affect different pathological processes of CAA, thus altering CAA-specific neuroimaging biomarkers. CAA has two pathological subtypes: CAA type 1 (parenchymal-predominant), characterised by amyloid in cortical capillaries and associated with APOE £4, and CAA type 2 (leptomeningealpredominant), where amyloid is deposited mainly in the leptomeningeal and cortical arteries and related to APOE ɛ2.^{26 27} A study revealed that FLP and SAE were significantly and specifically associated with cerebral microbleeds and cSS on MRI, respectively, which indicates that FLP and SAE may be related to the different anatomical distributions of CAA pathology: FLP may represent the parenchymal-predominant CAA (indicated by strictly lobar cerebral microbleeds), while SAE may represent the leptomeningeal-predominant CAA (indicated by cSS).^{24 28} A meta-analysis revealed a strong

SAE*	OR (95% CI)					
SAE*		P value	OR (95% CI)	P value	OR (95% CI)	P value
APOE ε2 dominar	nt					
Any allele	1.919 (1.107 to 3.325)	0.020	2.109 (1.167 to 3.810)	0.013	2.437 (1.218 to 4.875)	0.012
APOE ε 2 additive						
1 allele	1.917 (1.096 to 3.356)	0.067	2.131 (1.167 to 3.892)	0.046	2.547 (1.262 to 5.142)	0.032
2 alleles	1.948 (0.200 to 19.018)		1.708 (0.146 to 19.968)		0.774 (0.023 to 25.892)	
APOE ε4 dominar	nt					
Any allele	1.529 (0.864 to 2.705)	0.145	1.605 (0.873 to 2.950)	0.128	1.476 (0.728 to 2.996)	0.281
APOE <i>ɛ</i> 4 additive						
1 allele	1.612 (0.897 to 2.895)	0.267	1.677 (0.897 to 3.135)	0.266	1.627 (0.790 to 3.349)	0.292
2 alleles	0.793 (0.096 to 6.551)		0.917 (0.101 to 8.307)		0.238 (0.007 to 8.183)	
<i>APOE</i> ε2/ε4	6.744 (1.773 to 25.652)	0.005	7.599 (1.764 to 32.734)	0.006	8.855 (1.641 to 47.782)	0.011
FLP*						
APOE ε2 dominar	nt					
Any allele	1.302 (0.550 to 3.087)	0.548	1.722 (0.681 to 4.352)	0.251	1.732 (0.500 to 6.004)	0.386
APOE ε2 additive						
1 allele	1.364 (0.574 to 3.236)	0.781	1.795 (0.708 to 4.552)	0.469	1.770 (0.508 to 6.160)	0.669
2 alleles	-		-		-	
APOE ε4 dominar	nt					
Any allele	2.588 (1.241 to 5.395)	0.011	3.026 (1.353 to 6.767)	0.007	3.314 (1.128 to 9.732)	0.029
APOE ɛ4 additive						
1 allele	2.640 (1.237 to 5.636)	0.039	3.050 (1.328 to 7.006)	0.026	3.581 (1.198 to 10.705)	0.070
2 alleles	2.126 (0.252 to 17.96)		2.791 (0.274 to 28.419)		0.490 (0.001 to 181.012)	
<i>APOE</i> ε2/ε4	9.64 (2.073 to 44.834)	0.004	20.333 (3.278 to 126.137)	0.001	30.129 (1.832 to 495.633)	0.017
SAE+FLP*						
APOE ε2 dominar	nt					
Any allele	1.628 (0.675 to 3.93)	0.278	2.312 (0.881 to 6.073)	0.089	2.975 (0.77 to 11.502)	0.114
APOE ε 2 additive						
1 allele	1.704 (0.705 to 4.12)	0.496	2.408 (0.913 to 6.348)	0.206	3.023 (0.78 to 11.709)	0.278
2 alleles	-		-		-	
APOE ε4 dominar	nt					
Any allele	2.965 (1.36 to 6.464)	0.006	3.514 (1.485 to 8.316)	0.004	4.926 (1.425 to 17.028)	0.012
APOE ε4 additive						
1 allele	3 (1.341 to 6.712)	0.024	3.497 (1.435 to 8.522)	0.017	5.281 (1.502 to 18.57)	0.034
2 alleles	2.657 (0.312 to 22.653)		3.685 (0.344 to 39.458)		0.695 (0.001 to 805.727)	
APOE ε2/ε4	11.705 (2.497 to 54.871)	0.002	30.568 (4.460 to 209.503)	<0.001	70.360 (2.813 to 1759.792)	0.010

*Participants without either SAE or FLP were used as the comparator group.

APOE, apolipoprotein E; FLP, finger-like projections; ICH, intracerebral haemorrhage; SAE, subarachnoid extension.

association between *APOE* ε 2 genotype and cSS which is believed to be the chronic manifestation of episodes of SAE.^{18 20} Therefore, mechanistically, ε 2 may cause CAA of the leptomeningeal and cortical vessels and thus SAE. On the other hand, FLP may be a biomarker of more severe capillary CAA, which raises the likelihood of ICH dispersion into brain tissue, presumably driven by ε 4(24).

The strengths of this study include its use of a multicentre cohort and the use of standardised assessments of CAA-associated neuroimaging markers. This study analysed patients with supratentorial ICH rather than only lobar ICH, because the anatomical classification of ICH locations was not perfectly reliable, for example when an ICH is very large and extends into both lobar and non-lobar areas or multiple ICH including both areas. In addition, the analysis of $\epsilon 2/\epsilon 4$ heterozygosity showed that the association between $\epsilon 2/\epsilon 4$ and SAE was strong (OR=8.855), 3.6 times that of $\epsilon 2$ carrier status (OR=2.437), suggesting that the presence of the $\epsilon 4$ allele might augment the effect of ϵ 2. The association between ϵ 2/ ϵ 4 and FLP was strong (OR=30.129), 9.1 times that of ε 4 carrier status with FLP (OR=3.314), suggesting that ε_2 might increase the effect of £4. Analysis of the presence of FLP+SAE showed that its association with $\epsilon 2/\epsilon 4$ (OR=70.360) was 14.3 times that of $\varepsilon 4$ carrier statis (OR=4.926), which suggests that ε_2 and ε_4 independently affect SAE and FLP, and their joint effect further supports the hypothesis of a 'double hit' mechanism of $\epsilon 2/\epsilon 4$ heterozygosity.¹⁸ Based on the clinical significance of CT markers for CAA and the unidirectionality of genes in causal inference, the APOE genotype could be an important factor in the clinical evaluation of CAA diagnosis and prognosis, especially $\epsilon 2/\epsilon 4$ heterozygosity, which is likely to be a strong factor, although this genotype is rare.

Our study has limitations. First, the sample size of lobar ICH patients was relatively small, which may partly explain that the associations of APOE with imaging markers in these patients showed a same direction as all supratentorial ICH but not significant. Second, only CT axial plane was used to assess FLP because coronal and sagittal planes were not available, which may underestimate the prevalence of FLP. Third, given that the frequencies of APOE alleles varied by race and ethnicity,14 15 and the patients included in our study were from north China, the results should be interpreted with caution and need to be tested and verified in larger cohorts in future. On the other hand, similar prevalence of CAA neuropathology in (lobar) ICH patients in East-Asian and Western countries²⁹ might support the generalizability of our findings.

CONCLUSION

This study revealed that the $APOE \varepsilon 2$ and $\varepsilon 4$ alleles were associated with CT-based CAA neuroimaging markers (ie, FLP and SAE) in supratentorial ICH patients, which may indicate that different *APOE* alleles have different effects on the neuroimaging biomarkers and the potential pathophysiological processes.

Author affiliations

¹Department of Neurology, Peking University Third Hospital, Beijing, China ²Department of Radiology, Peking University Third Hospital, Beijing, China ³Department of Neurosurgery, Peking University Shougang Hospital, Beijing, China ⁴Research Center of Clincial Epidemiology, Peking University Third Hospital, Beijing, China

⁵Beijing Key Laboratory of Biomarker and Translational Research in

Neurodegenerative Diseases, China, Beijing, China

⁶Key Laboratory for Neuroscience, National Health Commission/Ministry of Education, Peking University, Beijing, China

Acknowledgements We are grateful to the patients and investigators in every centre for their involvement.

Contributors Conceptualisation, QY and DF; data curation, QY and LT; investigation, QY, XZ, XL, LT, FG and XH; methodology, QY and NL; formal analysis, QY and NL; supervision, DF; funding acquisition, QY and DF; writing—original draft, QY and XZ; writing—review and editing, LT, KX, NL and DF. All authors have read and agreed to the submitted version of the manuscript. DF is the guarantor.

rant number 910012

Funding National Natural Science Foundation of China (grant number 81901204) and Beijing Municipal Science and Technology Commission (grant number D141100000114005).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. This study was approved by the Ethics Committee of Peking University Third Hospital ((2014)-191-3). Participants gave informed consent to participate in the study before taking part. Provenance and peer review Not commissioned; externally peer reviewed. Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iD

Dongsheng Fan http://orcid.org/0000-0002-3129-9821

REFERENCES

- 1 Baron J-C, Jensen-Kondering U, Sacco S, *et al.* Can novel CTand MR-based neuroimaging biomarkers further improve the etiological diagnosis of lobar intra-cerebral hemorrhage? *J Neurol* 2023;270:582–8.
- 2 Sheth KN. Spontaneous Intracerebral Hemorrhage. N Engl J Med 2022;387:1589–96.
- 3 Greenberg SM, Ziai WC, Cordonnier C, et al. 2022 Guideline for the Management of Patients With Spontaneous Intracerebral Hemorrhage: A Guideline From the American Heart Association/ American Stroke Association. Stroke 2022;53:e282–361.
- 4 Samarasekera N, Rodrigues MA, Toh PS, et al. Imaging features of intracerebral hemorrhage with cerebral amyloid angiopathy: Systematic review and meta-analysis. PLoS One 2017;12:e0180923.
- 5 Rodrigues MA, Samarasekera N, Lerpiniere C, et al. The Edinburgh CT and genetic diagnostic criteria for lobar intracerebral haemorrhage associated with cerebral amyloid angiopathy: model development and diagnostic test accuracy study. *Lancet Neurol* 2018;17:232–40.
- 6 Sembill JA, Knott M, Xu M, et al. Simplified Edinburgh CT Criteria for Identification of Lobar Intracerebral Hemorrhage Associated With Cerebral Amyloid Angiopathy. *Neurology (ECronicon)* 2022;98:e1997–2004.
- 7 Greenberg SM. CT Diagnosis of Cerebral Amyloid Angiopathy: Blood Will Out. *Neurology (ECronicon)* 2022;98:823–4.
- 8 Raposo N, Charidimou A, Roongpiboonsopit D, *et al.* Convexity subarachnoid hemorrhage in lobar intracerebral hemorrhage: A prognostic marker. *Neurology (ECronicon)* 2020;94:e968–77.
- 9 Wilson D, Hostettler IC, Ambler G, et al. Convexity subarachnoid haemorrhage has a high risk of intracerebral haemorrhage in suspected cerebral amyloid angiopathy. J Neurol 2017;264:664–73.
- 10 Li Q, Zanon Zotin MC, Warren AD, et al. CT-Visible Convexity Subarachnoid Hemorrhage is Associated With Cortical Superficial Siderosis and Predicts Recurrent ICH. *Neurology (ECronicon)* 2021;96:e986–94.
- 11 Smith EE, Maas MB. To predict recurrence in cerebral amyloid angiopathy, look to the subarachnoid space. *Neurology (ECronicon)* 2020;94:375–6.
- 12 Ornello R, Colangeli E, Tommasino E, et al. Clinical usefulness of Edinburgh CT criteria in patients with lobar intracerebral hemorrhage. *Eur Stroke J* 2021;6:36–43.
- 13 Biffi A, Anderson CD, Jagiella JM, et al. APOE genotype and extent of bleeding and outcome in lobar intracerebral haemorrhage: a genetic association study. Lancet Neurol 2011;10:702–9.

- 14 Marini S, Crawford K, Morotti A, et al. Association of Apolipoprotein E With Intracerebral Hemorrhage Risk by Race/Ethnicity: A Metaanalysis. JAMA Neurol 2019;76:480–91.
- 15 Sawyer RP, Sekar P, Osborne J, et al. Racial/ethnic variation of APOE alleles for lobar intracerebral hemorrhage. *Neurology (ECronicon)* 2018;91:e410–20.
- 16 Rannikmäe K, Samarasekera N, Martînez-Gonzâlez NA, et al. Genetics of cerebral amyloid angiopathy: systematic review and meta-analysis. J Neurol Neurosurg Psychiatry 2013;84:901–8.
- 17 Rannikmäe K, Kalaria RN, Greenberg SM, et al. APOE associations with severe CAA-associated vasculopathic changes: collaborative meta-analysis. J Neurol Neurosurg Psychiatry 2014;85:300–5.
- 18 Charidimou A, Zonneveld HI, Shams S, et al. APOE and cortical superficial siderosis in CAA: Meta-analysis and potential mechanisms. *Neurology (ECronicon)* 2019;93:e358–71.
- 19 Greenberg SM, Rebeck GW, Vonsattel JP, et al. Apolipoprotein E epsilon 4 and cerebral hemorrhage associated with amyloid angiopathy. Ann Neurol 1995;38:254–9.
- 20 Charidimou A, Perosa V, Frosch MP, *et al*. Neuropathological correlates of cortical superficial siderosis in cerebral amyloid angiopathy. *Brain (Bacau)* 2020;143:3343–51.
- 21 Greenberg SM, Smith EE. Implications of cortical superficial siderosis in CAA: Superficial relationships. *Neurology (ECronicon)* 2019;92:360–1.

- 22 Mehndiratta P, Mendel TA. Cortical superficial siderosis, APOE genotype, and hemorrhage risk in cerebral amyloid angiopathy. *Neurol (ECronicon)* 2015;84:1190–1.
- 23 Ye X, Li G, Liu X, *et al.* Apolipoprotein E genotype predicts subarachnoid extension in spontaneous intracerebral haemorrhage. *Eur J Neurol* 2021;28:1992–9.
- 24 Hostettler IC, Seiffge D, Wong A, et al. APOE and Cerebral Small Vessel Disease Markers in Patients With Intracerebral Hemorrhage. Neurology (ECronicon) 2022;99:e1290–8.
- 25 Charidimou A, Schmitt A, Wilson D, et al. The Cerebral Haemorrhage Anatomical RaTing inStrument (CHARTS): Development and assessment of reliability. J Neurol Sci 2017;372:178–83.
- 26 Charidimou A, Boulouis G, Gurol ME, et al. Emerging concepts in sporadic cerebral amyloid angiopathy. *Brain (Bacau)* 2017;140:1829–50.
- 27 Thal DR, Ghebremedhin E, Rüb U, et al. Two types of sporadic cerebral amyloid angiopathy. J Neuropathol Exp Neurol 2002;61:282–93.
- 28 Schwarz G, Banerjee G, Hostettler IC, *et al.* MRI and CT imaging biomarkers of cerebral amyloid angiopathy in lobar intracerebral hemorrhage. *Int J Stroke* 2023;18:85–94.
- 29 De Kort AM, Verbeek MM, Schreuder FHBM, et al. Prevalence of Cerebral Amyloid Angiopathy Pathology and Strictly Lobar Microbleeds in East-Asian Versus Western Populations: A Systematic Review and Meta-Analysis. J Stroke 2024;26:179–89.