

Supplementary Table 5 Safety and efficacy endpoint comparison between different occlusion sites in PCO group by multivariable logistic regression.

Parameter	Recanalization		Any ICH within 24h		Symptomatic ICH within 24h	
	Adjust OR (95%CI)	p value	Adjust OR (95%CI)	p value	Adjust OR (95%CI)	p value
Occlusion site						
VA vs PCA	3.826 (1.201,12.190)	0.0232	1.239 (0.410,3.745)	0.7043	1.044 (0.086,12.658)	0.9728
BA vs PCA	3.748 (1.442,9.741)	0.0067	1.585 (0.592,4.237)	0.3597	1.068 (0.123,9.259)	0.9525
Age	1.003 (0.976,1.031)	0.8260	1.011 (0.988,1.036)	0.3532	1.051 (0.992,1.112)	0.0904
Male	1.073 (0.514,2.241)	0.8512	0.976 (0.514,1.855)	0.9412	0.790 (0.218,2.862)	0.7202
NIHSS	1.010 (0.982,1.039)	0.4701	1.002 (0.979,1.026)	0.8396	0.958 (0.908,1.011)	0.1215
Parameter	mRS 0-2 at 90d		mRS 0-3 at 90d		Mortality at 90d	
	Adjust OR (95%CI)	p value	Adjust OR (95%CI)	p value	Adjust OR (95%CI)	p value
Occlusion site						
VA vs PCA	0.777 (0.295,2.188)	0.6095	0.519 (0.219,1.394)	0.2106	NA	0.9446
BA vs PCA	0.921 (0.387,4.237)	0.8514	0.553 (0.236,2.584)	0.2094	NA	0.9460
Age	0.995 (0.977,1.014)	0.6295	1.002 (0.983,1.021)	0.8395	0.997 (0.974,1.021)	0.8142
Male	1.044 (0.615,1.771)	0.8744	1.407 (0.824,2.403)	0.2104	0.639 (0.338,1.206)	0.1667
NIHSS	0.947 (0.929,0.996)	<0.0001	0.942 (0.924,0.961)	<0.0001	1.059 (1.033,1.086)	<0.0001

VA, Vertebral artery; BA, Basilar artery; PCA, Posterior cerebral artery; ICH, intracranial hemorrhage; mRS, modified Rankin Scale; OR, odds ratio; NIHSS, National Institute of Health Stroke Scale.