

Cerebral hemodynamics in symptomatic intracranial atherosclerotic disease: A narrative review of the assessment methods and clinical implications

APPENDIX

Most of the images in the Figure in the main text were reproduced from previous publications with permissions, with the references provided below: image of positron emission tomography (PET),¹ single-photon emission CT (SPECT),² Xenon-enhanced CT (Xenon CT),³ carotid duplex ultrasound,⁴ CT perfusion (CTP),⁵ perfusion-weighted MR imaging (PWI),⁶ arterial spin labeling (ASL) MR perfusion,⁷ time-of-flight MR angiography (TOF-MRA),⁸ quantitative MR angiography or 4D flow MRI,⁹ digital subtraction angiography (DSA),¹⁰ and computational fluid dynamics (CFD) modeling.¹¹

REFERENCES

1. Grüner JM, Paamand R, Højgaard L, *et al.* Brain perfusion CT compared with ¹⁵O-H₂O-PET in healthy subjects. *EJNMMI Res* 2011;1:28.
2. Matsumoto Y, Ogasawara K, Saito H, *et al.* Detection of misery perfusion in the cerebral hemisphere with chronic unilateral major cerebral artery steno-occlusive disease using crossed cerebellar hypoperfusion: comparison of brain SPECT and PET imaging. *Eur J Nucl Med Mol Imaging* 2013;40:1573–81.
3. Rostami E, Engquist H, Howells T, *et al.* The correlation between cerebral blood flow measured by bedside Xenon-CT and brain chemistry monitored by microdialysis in the acute phase following subarachnoid hemorrhage. *Front Neurol* 2017;8:369.
4. Kim H, Byun E, Jeong MJ, *et al.* Duplex ultrasound findings and clinical outcomes of carotid restenosis after carotid endarterectomy. *PLoS One* 2020;15:e0244544.
5. Lan L, Leng X, Ip V, *et al.* Prolonged perfusion predicts recurrent ischemic stroke but not transient ischemic attack in patients with symptomatic intracranial stenosis. *Curr Neurovasc Res* 2017;14:149–57.
6. Bunevicius A, Yuan H, Lin W. The potential roles of ¹⁸F-FDG-PET in management of acute stroke patients. *Biomed Res Int* 2013;2013:634598.
7. Thamm T, Zweynert S, Piper SK, *et al.* Diagnostic and prognostic benefit of arterial spin labeling in subacute stroke. *Brain Behav* 2019;9:e01271.
8. Leng X, Wong KS, Soo Y, *et al.* Magnetic resonance angiography signal intensity as a

- marker of hemodynamic impairment in intracranial arterial stenosis. *PLoS One* 2013;8:e80124.
9. Miller KB, Howery AJ, Rivera-Rivera LA, *et al.* Age-related reductions in cerebrovascular reactivity using 4D flow MRI. *Front Aging Neurosci* 2019;11:281.
 10. Li L, Yang B, Dmytriw AA, *et al.* Hemodynamic versus anatomic assessment of symptomatic atherosclerotic middle cerebral artery stenosis: the relationship between pressure wire translesional gradient and angiographic lesion geometry. *Front Neurol* 2021;12:671778.
 11. Leng X, Lan L, Ip HL, *et al.* Translesional pressure gradient and leptomeningeal collateral status in symptomatic middle cerebral artery stenosis. *Eur J Neurol* 2018;25:404–10.